

On the Many Sorted Closure Operator and the Many Sorted Closure System

Artur Korniłowicz
Institute of Mathematics
Warsaw University
Białystok

MML Identifier: CLOSURE1.

WWW: <http://mizar.org/JFM/Vol8/closure1.html>

The articles [13], [4], [16], [12], [17], [2], [3], [6], [5], [14], [15], [1], [9], [10], [11], [8], and [7] provide the notation and terminology for this paper.

1. PRELIMINARIES

For simplicity, we adopt the following rules: i, x, I denote sets, A, M denote many sorted sets indexed by I , f denotes a function, and F denotes a many sorted function indexed by I .

The scheme *MSSUBSET* deals with a set \mathcal{A} , a non-empty many sorted set \mathcal{B} indexed by \mathcal{A} , a many sorted set \mathcal{C} indexed by \mathcal{A} , and a unary predicate \mathcal{P} , and states that:

If for every many sorted set X indexed by \mathcal{A} holds $X \in \mathcal{B}$ iff $X \in \mathcal{C}$ and $\mathcal{P}[X]$, then
 $\mathcal{B} \subseteq \mathcal{C}$

for all values of the parameters.

One can prove the following two propositions:

- (1) Let X be a non empty set and x, y be sets. If $x \subseteq y$, then $\{t; t \text{ ranges over elements of } X: y \subseteq t\} \subseteq \{z; z \text{ ranges over elements of } X: x \subseteq z\}$.
- (2) If there exists A such that $A \in M$, then M is non-empty.

Let us consider I, F, A . Then $F \leftarrow A$ is a many sorted set indexed by I .

Let us consider I , let A, B be non-empty many sorted sets indexed by I , let F be a many sorted function from A into B , and let X be an element of A . Then $F \leftarrow X$ is an element of B .

The following propositions are true:

- (3) Let A, X be many sorted sets indexed by I , B be a non-empty many sorted set indexed by I , and F be a many sorted function from A into B . If $X \in A$, then $F \leftarrow X \in B$.
- (4) Let F, G be many sorted functions indexed by I and A be a many sorted set indexed by I . If $A \in \text{dom}_\kappa G(\kappa)$, then $F \leftarrow (G \leftarrow A) = (F \circ G) \leftarrow A$.
- (5) If F is "1-1", then for all many sorted sets A, B indexed by I such that $A \in \text{dom}_\kappa F(\kappa)$ and $B \in \text{dom}_\kappa F(\kappa)$ and $F \leftarrow A = F \leftarrow B$ holds $A = B$.
- (6) Suppose $\text{dom}_\kappa F(\kappa)$ is non-empty and for all many sorted sets A, B indexed by I such that $A \in \text{dom}_\kappa F(\kappa)$ and $B \in \text{dom}_\kappa F(\kappa)$ and $F \leftarrow A = F \leftarrow B$ holds $A = B$. Then F is "1-1".

- (7) Let A, B be non-empty many sorted sets indexed by I and F, G be many sorted functions from A into B . If for every M such that $M \in A$ holds $F \leftarrow P M = G \leftarrow P M$, then $F = G$.

Let us consider I, M . Observe that there exists an element of 2^M which is empty yielding and locally-finite.

2. PROPERTIES OF MANY SORTED CLOSURE OPERATORS

Let us consider I, M . A set many sorted operation in M is a many sorted function from 2^M into 2^M .

Let us consider I, M , let O be a set many sorted operation in M , and let X be an element of 2^M . Then $O \leftarrow P X$ is an element of 2^M .

Let us consider I, M and let I_1 be a set many sorted operation in M . We say that I_1 is reflexive if and only if:

(Def. 2)¹ For every element X of 2^M holds $X \subseteq I_1 \leftarrow P X$.

We say that I_1 is monotonic if and only if:

(Def. 3) For all elements X, Y of 2^M such that $X \subseteq Y$ holds $I_1 \leftarrow P X \subseteq I_1 \leftarrow P Y$.

We say that I_1 is idempotent if and only if:

(Def. 4) For every element X of 2^M holds $I_1 \leftarrow P X = I_1 \leftarrow P (I_1 \leftarrow P X)$.

We say that I_1 is topological if and only if:

(Def. 5) For all elements X, Y of 2^M holds $I_1 \leftarrow P (X \cup Y) = I_1 \leftarrow P X \cup I_1 \leftarrow P Y$.

The following propositions are true:

(8) For every non-empty many sorted set M indexed by I and for every element X of M holds $X = \text{id}_M \leftarrow P X$.

(9) Let M be a non-empty many sorted set indexed by I and X, Y be elements of M . If $X \subseteq Y$, then $\text{id}_M \leftarrow P X \subseteq \text{id}_M \leftarrow P Y$.

(10) Let M be a non-empty many sorted set indexed by I and X, Y be elements of M . If $X \cup Y$ is an element of M , then $\text{id}_M \leftarrow P (X \cup Y) = \text{id}_M \leftarrow P X \cup \text{id}_M \leftarrow P Y$.

(11) Let X be an element of 2^M and i, x be sets. Suppose $i \in I$ and $x \in (\text{id}_{2^M} \leftarrow P X)(i)$. Then there exists a locally-finite element Y of 2^M such that $Y \subseteq X$ and $x \in (\text{id}_{2^M} \leftarrow P Y)(i)$.

Let us consider I, M . Observe that there exists a set many sorted operation in M which is reflexive, monotonic, idempotent, and topological.

The following propositions are true:

(12) id_{2^A} is a reflexive set many sorted operation in A .

(13) id_{2^A} is a monotonic set many sorted operation in A .

(14) id_{2^A} is an idempotent set many sorted operation in A .

(15) id_{2^A} is a topological set many sorted operation in A .

In the sequel P, R are set many sorted operations in M and E, T are elements of 2^M .

One can prove the following three propositions:

(16) If $E = M$ and P is reflexive, then $E = P \leftarrow P E$.

(17) If P is reflexive and for every element X of 2^M holds $P \leftarrow P X \subseteq X$, then P is idempotent.

¹ The definition (Def. 1) has been removed.

(18) If P is monotonic, then $P \leftrightarrow (E \cap T) \subseteq P \leftrightarrow E \cap P \leftrightarrow T$.

Let us consider I, M . One can verify that every set many sorted operation in M which is topological is also monotonic.

We now state the proposition

(19) If P is topological, then $P \leftrightarrow E \setminus P \leftrightarrow T \subseteq P \leftrightarrow (E \setminus T)$.

Let us consider I, M, R, P . Then $P \circ R$ is a set many sorted operation in M .

We now state several propositions:

(20) If P is reflexive and R is reflexive, then $P \circ R$ is reflexive.

(21) If P is monotonic and R is monotonic, then $P \circ R$ is monotonic.

(22) If P is idempotent and R is idempotent and $P \circ R = R \circ P$, then $P \circ R$ is idempotent.

(23) If P is topological and R is topological, then $P \circ R$ is topological.

(24) If P is reflexive and $i \in I$ and $f = P(i)$, then for every element x of $2^{M(i)}$ holds $x \subseteq f(x)$.

(25) If P is monotonic and $i \in I$ and $f = P(i)$, then for all elements x, y of $2^{M(i)}$ such that $x \subseteq y$ holds $f(x) \subseteq f(y)$.

(26) If P is idempotent and $i \in I$ and $f = P(i)$, then for every element x of $2^{M(i)}$ holds $f(x) = f(f(x))$.

(27) If P is topological and $i \in I$ and $f = P(i)$, then for all elements x, y of $2^{M(i)}$ holds $f(x \cup y) = f(x) \cup f(y)$.

3. ON THE MANY SORTED CLOSURE OPERATOR AND THE MANY SORTED CLOSURE SYSTEM

In the sequel S is a 1-sorted structure.

Let us consider S . We introduce many sorted closure system structures over S which are extensions of many-sorted structure over S and are systems

$\langle \text{sorts, a family} \rangle$,

where the sorts constitute a many sorted set indexed by the carrier of S and the family is a subset family of the sorts.

In the sequel M_1 is a many-sorted structure over S .

Let us consider S and let I_1 be a many sorted closure system structure over S . We say that I_1 is additive if and only if:

(Def. 6) The family of I_1 is additive.

We say that I_1 is absolutely-additive if and only if:

(Def. 7) The family of I_1 is absolutely-additive.

We say that I_1 is multiplicative if and only if:

(Def. 8) The family of I_1 is multiplicative.

We say that I_1 is absolutely-multiplicative if and only if:

(Def. 9) The family of I_1 is absolutely-multiplicative.

We say that I_1 is properly upper bound if and only if:

(Def. 10) The family of I_1 is properly upper bound.

We say that I_1 is properly lower bound if and only if:

(Def. 11) The family of I_1 is properly lower bound.

Let us consider S, M_1 . The functor $\text{MSFull}(M_1)$ yielding a many sorted closure system structure over S is defined as follows:

(Def. 12) $\text{MSFull}(M_1) = \langle \text{the sorts of } M_1, 2^{\text{the sorts of } M_1} \rangle$.

Let us consider S, M_1 . Note that $\text{MSFull}(M_1)$ is strict, additive, absolutely-additive, multiplicative, absolutely-multiplicative, properly upper bound, and properly lower bound.

Let us consider S and let M_1 be a non-empty many-sorted structure over S . One can verify that $\text{MSFull}(M_1)$ is non-empty.

Let us consider S . One can verify that there exists a many sorted closure system structure over S which is strict, non-empty, additive, absolutely-additive, multiplicative, absolutely-multiplicative, properly upper bound, and properly lower bound.

Let us consider S and let C_1 be an additive many sorted closure system structure over S . Observe that the family of C_1 is additive.

Let us consider S and let C_1 be an absolutely-additive many sorted closure system structure over S . Observe that the family of C_1 is absolutely-additive.

Let us consider S and let C_1 be a multiplicative many sorted closure system structure over S . One can verify that the family of C_1 is multiplicative.

Let us consider S and let C_1 be an absolutely-multiplicative many sorted closure system structure over S . Observe that the family of C_1 is absolutely-multiplicative.

Let us consider S and let C_1 be a properly upper bound many sorted closure system structure over S . One can check that the family of C_1 is properly upper bound.

Let us consider S and let C_1 be a properly lower bound many sorted closure system structure over S . Observe that the family of C_1 is properly lower bound.

Let us consider S , let M be a non-empty many sorted set indexed by the carrier of S , and let F be a subset family of M . Note that $\langle M, F \rangle$ is non-empty.

Let us consider S, M_1 and let F be an additive subset family of the sorts of M_1 . Note that $\langle \text{the sorts of } M_1, F \rangle$ is additive.

Let us consider S, M_1 and let F be an absolutely-additive subset family of the sorts of M_1 . Note that $\langle \text{the sorts of } M_1, F \rangle$ is absolutely-additive.

Let us consider S, M_1 and let F be a multiplicative subset family of the sorts of M_1 . Note that $\langle \text{the sorts of } M_1, F \rangle$ is multiplicative.

Let us consider S, M_1 and let F be an absolutely-multiplicative subset family of the sorts of M_1 . Observe that $\langle \text{the sorts of } M_1, F \rangle$ is absolutely-multiplicative.

Let us consider S, M_1 and let F be a properly upper bound subset family of the sorts of M_1 . Note that $\langle \text{the sorts of } M_1, F \rangle$ is properly upper bound.

Let us consider S, M_1 and let F be a properly lower bound subset family of the sorts of M_1 . Observe that $\langle \text{the sorts of } M_1, F \rangle$ is properly lower bound.

Let us consider S . Observe that every many sorted closure system structure over S which is absolutely-additive is also additive.

Let us consider S . Observe that every many sorted closure system structure over S which is absolutely-multiplicative is also multiplicative.

Let us consider S . One can verify that every many sorted closure system structure over S which is absolutely-multiplicative is also properly upper bound.

Let us consider S . Note that every many sorted closure system structure over S which is absolutely-additive is also properly lower bound.

Let us consider S . A many sorted closure system of S is an absolutely-multiplicative many sorted closure system structure over S .

Let us consider I, M . A many sorted closure operator of M is a reflexive monotonic idempotent set many sorted operation in M .

Let us consider I, M and let F be a many sorted function from M into M . The functor $\text{FixPoints}(F)$ yields a many sorted subset indexed by M and is defined as follows:

(Def. 13) For every i such that $i \in I$ holds $x \in (\text{FixPoints}(F))(i)$ iff there exists a function f such that $f = F(i)$ and $x \in \text{dom } f$ and $f(x) = x$.

Let us consider I , let M be an empty yielding many sorted set indexed by I , and let F be a many sorted function from M into M . Note that $\text{FixPoints}(F)$ is empty yielding.

Next we state a number of propositions:

- (28) For every many sorted function F from M into M holds $A \in M$ and $F \leftrightarrow A = A$ iff $A \in \text{FixPoints}(F)$.
- (29) $\text{FixPoints}(\text{id}_A) = A$.
- (30) Let A be a many sorted set indexed by the carrier of S , J be a reflexive monotonic set many sorted operation in A , and D be a subset family of A . If $D = \text{FixPoints}(J)$, then $\langle A, D \rangle$ is a many sorted closure system of S .
- (31) Let D be a properly upper bound subset family of M and X be an element of 2^M . Then there exists a non-empty subset family S_1 of M such that for every many sorted set Y indexed by I holds $Y \in S_1$ if and only if the following conditions are satisfied:
- (i) $Y \in D$, and
 - (ii) $X \subseteq Y$.
- (32) Let D be a properly upper bound subset family of M , X be an element of 2^M , and S_1 be a non-empty subset family of M . Suppose that for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$. Let i be a set and D_1 be a non empty set. If $i \in I$ and $D_1 = D(i)$, then $S_1(i) = \{z; z \text{ ranges over elements of } D_1: X(i) \subseteq z\}$.
- (33) Let D be a properly upper bound subset family of M . Then there exists a set many sorted operation J in M such that for every element X of 2^M and for every non-empty subset family S_1 of M if for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$, then $J \leftrightarrow X = \bigcap S_1$.
- (34) Let D be a properly upper bound subset family of M , A be an element of 2^M , and J be a set many sorted operation in M . Suppose that
- (i) $A \in D$, and
 - (ii) for every element X of 2^M and for every non-empty subset family S_1 of M such that for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$ holds $J \leftrightarrow X = \bigcap S_1$. Then $J \leftrightarrow A = A$.
- (35) Let D be an absolutely-multiplicative subset family of M , A be an element of 2^M , and J be a set many sorted operation in M . Suppose that
- (i) $J \leftrightarrow A = A$, and
 - (ii) for every element X of 2^M and for every non-empty subset family S_1 of M such that for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$ holds $J \leftrightarrow X = \bigcap S_1$. Then $A \in D$.
- (36) Let D be a properly upper bound subset family of M and J be a set many sorted operation in M . Suppose that for every element X of 2^M and for every non-empty subset family S_1 of M such that for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$ holds $J \leftrightarrow X = \bigcap S_1$. Then J is reflexive and monotonic.
- (37) Let D be an absolutely-multiplicative subset family of M and J be a set many sorted operation in M . Suppose that for every element X of 2^M and for every non-empty subset family S_1 of M such that for every many sorted set Y indexed by I holds $Y \in S_1$ iff $Y \in D$ and $X \subseteq Y$ holds $J \leftrightarrow X = \bigcap S_1$. Then J is idempotent.
- (38) Let D be a many sorted closure system of S and J be a set many sorted operation in the sorts of D . Suppose that for every element X of $2^{\text{the sorts of } D}$ and for every non-empty subset family S_1 of the sorts of D such that for every many sorted set Y indexed by the carrier of S holds $Y \in S_1$ iff $Y \in$ the family of D and $X \subseteq Y$ holds $J \leftrightarrow X = \bigcap S_1$. Then J is a many sorted closure operator of the sorts of D .

Let us consider S , let A be a many sorted set indexed by the carrier of S , and let C be a many sorted closure operator of A . The functor $\text{ClSys}(C)$ yields a many sorted closure system of S and is defined by:

(Def. 14) There exists a subset family D of A such that $D = \text{FixPoints}(C)$ and $\text{ClSys}(C) = \langle A, D \rangle$.

Let us consider S , let A be a many sorted set indexed by the carrier of S , and let C be a many sorted closure operator of A . Observe that $\text{ClSys}(C)$ is strict.

Let us consider S , let A be a non-empty many sorted set indexed by the carrier of S , and let C be a many sorted closure operator of A . One can check that $\text{ClSys}(C)$ is non-empty.

Let us consider S and let D be a many sorted closure system of S . The functor $\text{ClOp}(D)$ yields a many sorted closure operator of the sorts of D and is defined by the condition (Def. 15).

(Def. 15) Let X be an element of $2^{\text{the sorts of } D}$ and S_1 be a non-empty subset family of the sorts of D . Suppose that for every many sorted set Y indexed by the carrier of S holds $Y \in S_1$ iff $Y \in$ the family of D and $X \subseteq Y$. Then $(\text{ClOp}(D)) \dashv\vdash X = \bigcap S_1$.

The following propositions are true:

(39) Let A be a many sorted set indexed by the carrier of S and J be a many sorted closure operator of A . Then $\text{ClOp}(\text{ClSys}(J)) = J$.

(40) For every many sorted closure system D of S holds $\text{ClSys}(\text{ClOp}(D)) =$ the many sorted closure system structure of D .

REFERENCES

- [1] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_2.html.
- [2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [4] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [5] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [6] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [7] Artur Kornilowicz. Certain facts about families of subsets of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/mssubfam.html>.
- [8] Artur Kornilowicz. Extensions of mappings on generator set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/extens_1.html.
- [9] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_3.html.
- [10] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [11] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pre_circ.html.
- [12] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [14] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [15] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [16] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

- [17] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received February 7, 1996

Published January 2, 2004
