

Algebraic and Arithmetic Lattices. Part I¹

Robert Milewski
Warsaw University
Białystok

Summary. We formalize [10, pp.87–89]

MML Identifier: WAYBEL13.

The papers [17], [21], [20], [16], [14], [9], [22], [19], [6], [7], [15], [18], [1], [2], [11], [24], [4], [8], [5], [23], [12], [3], and [13] provide the terminology and notation for this paper.

1. PRELIMINARIES

The scheme *LambdaCD* deals with a non empty set \mathcal{A} , a unary functor \mathcal{F} yielding a set, a unary functor \mathcal{G} yielding a set, and a unary predicate \mathcal{P} , and states that:

There exists a function f such that $\text{dom } f = \mathcal{A}$ and for every element x of \mathcal{A} holds if $\mathcal{P}[x]$, then $f(x) = \mathcal{F}(x)$ and if not $\mathcal{P}[x]$, then $f(x) = \mathcal{G}(x)$

for all values of the parameters.

The following propositions are true:

- (1) Let L be a non empty reflexive transitive relational structure and x, y be elements of L . If $x \leq y$, then $\text{compactbelow}(x) \subseteq \text{compactbelow}(y)$.
- (2) For every non empty reflexive relational structure L and for every element x of L holds $\text{compactbelow}(x)$ is a subset of $\text{CompactSublatt}(L)$.
- (3) For every relational structure L and for every relational substructure S of L holds every subset of S is a subset of L .

¹This work was partially supported by KBN Grant 8 T11C 018 12.

- (4) For every non empty reflexive transitive relational structure L with l.u.b.'s holds the carrier of L is an ideal of L .
- (5) Let L_1 be a lower-bounded non empty reflexive antisymmetric relational structure and L_2 be a non empty reflexive antisymmetric relational structure. Suppose the relational structure of $L_1 =$ the relational structure of L_2 and L_1 is up-complete. Then the carrier of $\text{CompactSublatt}(L_1) =$ the carrier of $\text{CompactSublatt}(L_2)$.

2. ALGEBRAIC AND ARITHMETIC LATTICES

Next we state three propositions:

- (6) For every algebraic lower-bounded lattice L holds every continuous subframe of L is algebraic.
- (7) Let X, E be sets and L be a continuous subframe of 2_{\subseteq}^X . Then $E \in$ the carrier of $\text{CompactSublatt}(L)$ if and only if there exists an element F of 2_{\subseteq}^X such that F is finite and $E = \bigcap \{Y, Y \text{ ranges over elements of } L: F \subseteq Y\}$ and $F \subseteq E$.
- (8) For every lower-bounded sup-semilattice L holds $\langle \text{Ids}(L), \subseteq \rangle$ is a continuous subframe of $2_{\subseteq}^{\text{the carrier of } L}$.

Let L be a non empty reflexive transitive relational structure. Observe that there exists an ideal of L which is principal.

One can prove the following propositions:

- (9) For every lower-bounded sup-semilattice L and for every non empty directed subset X of $\langle \text{Ids}(L), \subseteq \rangle$ holds $\text{sup } X = \bigcup X$.
- (10) For every lower-bounded sup-semilattice S holds $\langle \text{Ids}(S), \subseteq \rangle$ is algebraic.
- (11) Let S be a lower-bounded sup-semilattice and x be an element of $\langle \text{Ids}(S), \subseteq \rangle$. Then x is compact if and only if x is a principal ideal of S .
- (12) Let S be a lower-bounded sup-semilattice and x be an element of $\langle \text{Ids}(S), \subseteq \rangle$. Then x is compact if and only if there exists an element a of S such that $x = \downarrow a$.
- (13) Let L be a lower-bounded sup-semilattice and f be a map from L into $\text{CompactSublatt}(\langle \text{Ids}(L), \subseteq \rangle)$. If for every element x of L holds $f(x) = \downarrow x$, then f is isomorphic.
- (14) For every lower-bounded lattice S holds $\langle \text{Ids}(S), \subseteq \rangle$ is arithmetic.
- (15) For every lower-bounded sup-semilattice L holds $\text{CompactSublatt}(L)$ is a lower-bounded sup-semilattice.

- (16) Let L be an algebraic lower-bounded sup-semilattice and f be a map from L into $\langle \text{Ids}(\text{CompactSublatt}(L)), \subseteq \rangle$. If for every element x of L holds $f(x) = \text{compactbelow}(x)$, then f is isomorphic.
- (17) Let L be an algebraic lower-bounded sup-semilattice and x be an element of L . Then $\text{compactbelow}(x)$ is a principal ideal of $\text{CompactSublatt}(L)$ if and only if x is compact.

3. MAPS

We now state three propositions:

- (18) Let L_1, L_2 be non empty relational structures, X be a subset of L_1 , x be an element of L_1 , and f be a map from L_1 into L_2 . If f is isomorphic, then $x \leq X$ iff $f(x) \leq f^\circ X$.
- (19) Let L_1, L_2 be non empty relational structures, X be a subset of L_1 , x be an element of L_1 , and f be a map from L_1 into L_2 . If f is isomorphic, then $x \geq X$ iff $f(x) \geq f^\circ X$.
- (20) Let L_1, L_2 be non empty antisymmetric relational structures and f be a map from L_1 into L_2 . If f is isomorphic, then f is infs-preserving and sups-preserving.

Let L_1, L_2 be non empty antisymmetric relational structures. Note that every map from L_1 into L_2 which is isomorphic is also infs-preserving and sups-preserving.

We now state a number of propositions:

- (21) Let L_1, L_2, L_3 be non empty transitive antisymmetric relational structures and f be a map from L_1 into L_2 . Suppose f is infs-preserving. Suppose L_2 is a full infs-inheriting relational substructure of L_3 and L_3 is complete. Then there exists a map g from L_1 into L_3 such that $f = g$ and g is infs-preserving.
- (22) Let L_1, L_2, L_3 be non empty transitive antisymmetric relational structures and f be a map from L_1 into L_2 . Suppose f is monotone and directed-sups-preserving. Suppose L_2 is a full directed-sups-inheriting relational substructure of L_3 and L_3 is complete. Then there exists a map g from L_1 into L_3 such that $f = g$ and g is directed-sups-preserving.
- (23) For every lower-bounded sup-semilattice L holds $\langle \text{Ids}(\text{CompactSublatt}(L)), \subseteq \rangle$ is a continuous subframe of $2_{\subseteq}^{\text{the carrier of CompactSublatt}(L)}$.
- (24) Let L be an algebraic lower-bounded lattice. Then there exists a map g from L into $2_{\subseteq}^{\text{the carrier of CompactSublatt}(L)}$ such that
- (i) g is infs-preserving, directed-sups-preserving, and one-to-one, and
 - (ii) for every element x of L holds $g(x) = \text{compactbelow}(x)$.

- (25) Let I be a non empty set and J be a relational structure yielding nonempty reflexive-yielding many sorted set indexed by I . Suppose that for every element i of I holds $J(i)$ is an algebraic lower-bounded lattice. Then $\prod J$ is an algebraic lower-bounded lattice.
- (26) Let L_1, L_2 be non empty relational structures. Suppose the relational structure of $L_1 =$ the relational structure of L_2 . Then L_1 and L_2 are isomorphic.
- (27) Let L_1, L_2 be up-complete non empty posets and f be a map from L_1 into L_2 . Suppose f is isomorphic. Let x, y be elements of L_1 . Then $x \ll y$ if and only if $f(x) \ll f(y)$.
- (28) Let L_1, L_2 be up-complete non empty posets and f be a map from L_1 into L_2 . Suppose f is isomorphic. Let x be an element of L_1 . Then x is compact if and only if $f(x)$ is compact.
- (29) Let L_1, L_2 be up-complete non empty posets and f be a map from L_1 into L_2 . If f is isomorphic, then for every element x of L_1 holds $f \circ \text{compactbelow}(x) = \text{compactbelow}(f(x))$.
- (30) For all non empty posets L_1, L_2 such that L_1 and L_2 are isomorphic and L_1 is up-complete holds L_2 is up-complete.
- (31) For all non empty posets L_1, L_2 such that L_1 and L_2 are isomorphic and L_1 is complete and satisfies axiom K holds L_2 satisfies axiom K.
- (32) Let L_1, L_2 be sup-semilattices. Suppose L_1 and L_2 are isomorphic and L_1 is lower-bounded and algebraic. Then L_2 is algebraic.
- (33) For every continuous lower-bounded sup-semilattice L holds $\text{SupMap}(L)$ is infs-preserving and sups-preserving.
- (34) Let L be a lower-bounded lattice. Then L is algebraic if and only if there exists a set X and there exists a full relational substructure S of 2_{\subseteq}^X such that S is infs-inheriting and directed-sups-inheriting and L and S are isomorphic.
- (35) Let L be a lower-bounded lattice. Then L is algebraic if and only if there exists a set X and there exists a closure map c from 2_{\subseteq}^X into 2_{\subseteq}^X such that c is directed-sups-preserving and L and $\text{Im } c$ are isomorphic.

REFERENCES

- [1] Grzegorz Bancerek. Complete lattices. *Formalized Mathematics*, 2(5):719–725, 1991.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. *Formalized Mathematics*, 6(1):81–91, 1997.
- [3] Grzegorz Bancerek. Closure operators and subalgebras. *Formalized Mathematics*, 6(2):295–301, 1997.
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Formalized Mathematics*, 6(1):93–107, 1997.
- [5] Grzegorz Bancerek. The “way-below” relation. *Formalized Mathematics*, 6(1):169–176, 1997.

- [6] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [7] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [8] Czesław Byliński. Galois connections. *Formalized Mathematics*, 6(1):131–143, 1997.
- [9] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [10] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [11] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Formalized Mathematics*, 6(1):117–121, 1997.
- [12] Robert Milewski. Algebraic lattices. *Formalized Mathematics*, 6(2):249–254, 1997.
- [13] Michał Muzalewski. Categories of groups. *Formalized Mathematics*, 2(4):563–571, 1991.
- [14] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(1):147–152, 1990.
- [15] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [16] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [17] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [18] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(1):15–22, 1993.
- [19] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(2):313–319, 1990.
- [20] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [21] Zinaida Trybulec and Halina Świączkowska. Boolean properties of sets. *Formalized Mathematics*, 1(1):17–23, 1990.
- [22] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [23] Mariusz Żynel. The equational characterization of continuous lattices. *Formalized Mathematics*, 6(2):199–205, 1997.
- [24] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Formalized Mathematics*, 6(1):123–130, 1997.

Received March 4, 1997
