

Predicate Calculus for Boolean Valued Functions. Part X

Shunichi Kobayashi
Ueda Multimedia Information Center
Nagano

Summary. In this paper, we proved some elementary predicate calculus formulae containing the quantifiers of Boolean valued functions with respect to partitions. Such a theory is an analogy of usual predicate logic.

MML Identifier: BVFUNC18.

The notation and terminology used here are introduced in the following articles: [1], [2], [3], [4], and [5].

In this paper Y is a non empty set.

One can prove the following propositions:

- (1) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\neg \exists_{a,AG,B}G \in \exists_{\forall -a,B}G,AG$.
- (2) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\neg \exists_{a,AG,B}G \in \forall_{\forall -a,B}G,AG$.
- (3) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\exists_{\neg \exists a,AG,B}G \in \neg \exists_{\forall a,B}G,AG$.
- (4) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg \exists a,AG,B}G \in \neg \exists_{\forall a,B}G,AG$.
- (5) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg \exists a,AG,B}G \in \neg \forall_{\exists a,B}G,AG$.

- (6) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\exists_{a,A}G,B}G \in \neg\exists_{\exists_{a,B}G,A}G$.
- (7) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\exists_{\neg\forall_{a,A}G,B}G \in \exists_{\neg\forall_{a,B}G,A}G$.
- (8) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\forall_{a,A}G,B}G \in \exists_{\neg\forall_{a,B}G,A}G$.
- (9) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\exists_{\neg\exists_{a,A}G,B}G \in \exists_{\neg\forall_{a,B}G,A}G$.
- (10) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\exists_{a,A}G,B}G \in \exists_{\neg\forall_{a,B}G,A}G$.
- (11) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\exists_{\neg\exists_{a,A}G,B}G \in \forall_{\neg\forall_{a,B}G,A}G$.
- (12) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\exists_{a,A}G,B}G \in \forall_{\neg\forall_{a,B}G,A}G$.
- (13) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\exists_{a,A}G,B}G \in \exists_{\neg\exists_{a,B}G,A}G$.
- (14) Let a be an element of $BVF(Y)$, G be a subset of $PARTITIONS(Y)$, and A, B, C be partitions of Y . Suppose G is a coordinate and $G = \{A, B, C\}$ and $A \neq B$ and $B \neq C$ and $C \neq A$. Then $\forall_{\neg\exists_{a,A}G,B}G \in \forall_{\neg\exists_{a,B}G,A}G$.

REFERENCES

- [1] Shunichi Kobayashi and Kui Jia. A theory of Boolean valued functions and partitions. *Formalized Mathematics*, 7(2):249–254, 1998.
- [2] Shunichi Kobayashi and Yatsuka Nakamura. A theory of Boolean valued functions and quantifiers with respect to partitions. *Formalized Mathematics*, 7(2):307–312, 1998.
- [3] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Formalized Mathematics*, 1(3):441–444, 1990.
- [4] Andrzej Trybulec. Semilattice operations on finite subsets. *Formalized Mathematics*, 1(2):369–376, 1990.
- [5] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.

Received November 15, 1999
