

Some Properties of Cells on Go Board

Czesław Byliński
University of Białystok

MML Identifier: GOBRD13.

WWW: <http://mizar.org/JFM/Vol11/gobrd13.html>

The articles [20], [7], [22], [2], [18], [23], [5], [6], [1], [4], [8], [21], [9], [17], [3], [13], [12], [19], [10], [11], [14], [15], and [16] provide the notation and terminology for this paper.

We adopt the following convention: $i, i_1, i_2, j, j_1, j_2, k, n$ denote natural numbers, D denotes a non empty set, and f denotes a finite sequence of elements of D .

Let E be a non empty set, let S be a non empty set of finite sequences of the carrier of \mathcal{E}_T^2 , let F be a function from E into S , and let e be an element of E . Then $F(e)$ is a finite sequence of elements of \mathcal{E}_T^2 .

Let F be a function. The functor $\text{Values } F$ yields a set and is defined by:

(Def. 1) $\text{Values } F = \bigcup (\text{rng}_\kappa F(\kappa))$.

One can prove the following proposition

(1) For every finite sequence M of elements of D^* holds $M(i)$ is a finite sequence of elements of D .

Let D be a set. Observe that every finite sequence of elements of D^* is finite sequence yielding. One can check that every function which is finite sequence yielding is also function yielding.

One can prove the following proposition

(3)¹ For every finite sequence M of elements of D^* holds $\text{Values } M = \bigcup \{\text{rng } f; f \text{ ranges over elements of } D^*: f \in \text{rng } M\}$.

Let D be a non empty set and let M be a finite sequence of elements of D^* . One can check that $\text{Values } M$ is finite.

One can prove the following propositions:

(4) For every matrix M over D such that $i \in \text{dom } M$ and $M(i) = f$ holds $\text{len } f = \text{width } M$.

(5) For every matrix M over D such that $i \in \text{dom } M$ and $M(i) = f$ and $j \in \text{dom } f$ holds $\langle i, j \rangle \in \text{the indices of } M$.

(6) For every matrix M over D such that $\langle i, j \rangle \in \text{the indices of } M$ and $M(i) = f$ holds $\text{len } f = \text{width } M$ and $j \in \text{dom } f$.

(7) For every matrix M over D holds $\text{Values } M = \{M \circ \langle i, j \rangle : \langle i, j \rangle \in \text{the indices of } M\}$.

(8) For every non empty set D and for every matrix M over D holds $\text{card } \text{Values } M \leq \text{len } M \cdot \text{width } M$.

¹ The proposition (2) has been removed.

In the sequel f is a finite sequence of elements of \mathcal{E}_T^2 and G is a Go-board.
The following propositions are true:

- (9) For every matrix G over \mathcal{E}_T^2 such that f is a sequence which elements belong to G holds $\text{rng } f \subseteq \text{Values } G$.
- (10) For all Go-boards G_1, G_2 such that $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $1 \leq j_2$ and $j_2 \leq \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (1, j_2)$ holds $i_1 = 1$.
- (11) For all Go-boards G_1, G_2 such that $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $1 \leq j_2$ and $j_2 \leq \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (\text{len } G_2, j_2)$ holds $i_1 = \text{len } G_1$.
- (12) For all Go-boards G_1, G_2 such that $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $1 \leq i_2$ and $i_2 \leq \text{len } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, 1)$ holds $j_1 = 1$.
- (13) For all Go-boards G_1, G_2 such that $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $1 \leq i_2$ and $i_2 \leq \text{len } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, \text{width } G_2)$ holds $j_1 = \text{width } G_1$.
- (14) Let G_1, G_2 be Go-boards. Suppose $\text{Values } G_1 \subseteq \text{Values } G_2$ and $1 \leq i_1$ and $i_1 < \text{len } G_1$ and $1 \leq j_1$ and $j_1 \leq \text{width } G_1$ and $1 \leq i_2$ and $i_2 < \text{len } G_2$ and $1 \leq j_2$ and $j_2 \leq \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $(G_2 \circ (i_2 + 1, j_2))_1 \leq (G_1 \circ (i_1 + 1, j_1))_1$.
- (15) Let G_1, G_2 be Go-boards. Suppose $G_1 \circ (i_1 - 1, j_1) \in \text{Values } G_2$ and $1 < i_1$ and $i_1 \leq \text{len } G_1$ and $1 \leq j_1$ and $j_1 \leq \text{width } G_1$ and $1 < i_2$ and $i_2 \leq \text{len } G_2$ and $1 \leq j_2$ and $j_2 \leq \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $(G_1 \circ (i_1 - 1, j_1))_1 \leq (G_2 \circ (i_2 - 1, j_2))_1$.
- (16) Let G_1, G_2 be Go-boards. Suppose $G_1 \circ (i_1, j_1 + 1) \in \text{Values } G_2$ and $1 \leq i_1$ and $i_1 \leq \text{len } G_1$ and $1 \leq j_1$ and $j_1 < \text{width } G_1$ and $1 \leq i_2$ and $i_2 \leq \text{len } G_2$ and $1 \leq j_2$ and $j_2 < \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $(G_2 \circ (i_2, j_2 + 1))_2 \leq (G_1 \circ (i_1, j_1 + 1))_2$.
- (17) Let G_1, G_2 be Go-boards. Suppose $\text{Values } G_1 \subseteq \text{Values } G_2$ and $1 \leq i_1$ and $i_1 \leq \text{len } G_1$ and $1 < j_1$ and $j_1 \leq \text{width } G_1$ and $1 \leq i_2$ and $i_2 \leq \text{len } G_2$ and $1 < j_2$ and $j_2 \leq \text{width } G_2$ and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $(G_1 \circ (i_1, j_1 - 1))_2 \leq (G_2 \circ (i_2, j_2 - 1))_2$.
- (18) Let G_1, G_2 be Go-boards. Suppose $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $\langle i_2, j_2 \rangle \in$ the indices of G_2 and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $\text{cell}(G_2, i_2, j_2) \subseteq \text{cell}(G_1, i_1, j_1)$.
- (19) Let G_1, G_2 be Go-boards. Suppose $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $\langle i_2, j_2 \rangle \in$ the indices of G_2 and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $\text{cell}(G_2, i_2 - 1, j_2) \subseteq \text{cell}(G_1, i_1 - 1, j_1)$.
- (20) Let G_1, G_2 be Go-boards. Suppose $\text{Values } G_1 \subseteq \text{Values } G_2$ and $\langle i_1, j_1 \rangle \in$ the indices of G_1 and $\langle i_2, j_2 \rangle \in$ the indices of G_2 and $G_1 \circ (i_1, j_1) = G_2 \circ (i_2, j_2)$. Then $\text{cell}(G_2, i_2, j_2 - 1) \subseteq \text{cell}(G_1, i_1, j_1 - 1)$.
- (21) Let f be a standard special circular sequence. Suppose f is a sequence which elements belong to G . Then $\text{Values the Go-board of } f \subseteq \text{Values } G$.

Let us consider f, G, k . Let us assume that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G . The functor $\text{right_cell}(f, k, G)$ yields a subset of \mathcal{E}_T^2 and is defined by the condition (Def. 2).

- (Def. 2) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\text{right_cell}(f, k, G) = \text{cell}(G, i_1, j_1)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\text{right_cell}(f, k, G) = \text{cell}(G, i_1, j_1 - 1)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\text{right_cell}(f, k, G) = \text{cell}(G, i_2, j_2)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\text{right_cell}(f, k, G) = \text{cell}(G, i_1 - 1, j_2)$.

The functor $\text{left_cell}(f, k, G)$ yielding a subset of \mathcal{E}_T^2 is defined by the condition (Def. 3).

(Def. 3) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then

- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\text{left_cell}(f, k, G) = \text{cell}(G, i_1 -' 1, j_1)$, or
- (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\text{left_cell}(f, k, G) = \text{cell}(G, i_1, j_1)$, or
- (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\text{left_cell}(f, k, G) = \text{cell}(G, i_2, j_2 -' 1)$, or
- (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\text{left_cell}(f, k, G) = \text{cell}(G, i_1, j_2)$.

One can prove the following propositions:

- (22) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i, j + 1 \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i, j + 1)$. Then $\text{left_cell}(f, k, G) = \text{cell}(G, i -' 1, j)$.
- (23) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i, j + 1 \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i, j + 1)$. Then $\text{right_cell}(f, k, G) = \text{cell}(G, i, j)$.
- (24) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i + 1, j \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i + 1, j)$. Then $\text{left_cell}(f, k, G) = \text{cell}(G, i, j)$.
- (25) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i + 1, j \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i + 1, j)$. Then $\text{right_cell}(f, k, G) = \text{cell}(G, i, j -' 1)$.
- (26) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i + 1, j \rangle \in$ the indices of G and $f_k = G \circ (i + 1, j)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{left_cell}(f, k, G) = \text{cell}(G, i, j -' 1)$.
- (27) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i + 1, j \rangle \in$ the indices of G and $f_k = G \circ (i + 1, j)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{right_cell}(f, k, G) = \text{cell}(G, i, j)$.
- (28) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j + 1 \rangle \in$ the indices of G and $\langle i, j \rangle \in$ the indices of G and $f_k = G \circ (i, j + 1)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{left_cell}(f, k, G) = \text{cell}(G, i, j)$.
- (29) Suppose that $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j + 1 \rangle \in$ the indices of G and $\langle i, j \rangle \in$ the indices of G and $f_k = G \circ (i, j + 1)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{right_cell}(f, k, G) = \text{cell}(G, i -' 1, j)$.
- (30) If $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G , then $\text{left_cell}(f, k, G) \cap \text{right_cell}(f, k, G) = \mathcal{L}(f, k)$.
- (31) If $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G , then $\text{right_cell}(f, k, G)$ is closed.
- (32) Suppose $1 \leq k$ and $k + 1 \leq \text{len } f$ and f is a sequence which elements belong to G and $k + 1 \leq n$. Then $\text{left_cell}(f, k, G) = \text{left_cell}(f|_n, k, G)$ and $\text{right_cell}(f, k, G) = \text{right_cell}(f|_n, k, G)$.
- (33) Suppose $1 \leq k$ and $k + 1 \leq \text{len}(f|_n)$ and $n \leq \text{len } f$ and f is a sequence which elements belong to G . Then $\text{left_cell}(f, k + n, G) = \text{left_cell}(f|_n, k, G)$ and $\text{right_cell}(f, k + n, G) = \text{right_cell}(f|_n, k, G)$.
- (34) Let G be a Go-board and f be a standard special circular sequence. Suppose $1 \leq n$ and $n + 1 \leq \text{len } f$ and f is a sequence which elements belong to G . Then $\text{left_cell}(f, n, G) \subseteq \text{leftcell}(f, n)$ and $\text{right_cell}(f, n, G) \subseteq \text{rightcell}(f, n)$.

Let us consider f, G, k . Let us assume that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G . The functor $\text{front_right_cell}(f, k, G)$ yields a subset of \mathcal{E}_T^2 and is defined by the condition (Def. 4).

- (Def. 4) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\text{front_right_cell}(f, k, G) = \text{cell}(G, i_2, j_2)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\text{front_right_cell}(f, k, G) = \text{cell}(G, i_2, j_2 -' 1)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\text{front_right_cell}(f, k, G) = \text{cell}(G, i_2 -' 1, j_2)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\text{front_right_cell}(f, k, G) = \text{cell}(G, i_2 -' 1, j_2 -' 1)$.

The functor $\text{front_left_cell}(f, k, G)$ yielding a subset of \mathcal{E}_T^2 is defined by the condition (Def. 5).

- (Def. 5) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\text{front_left_cell}(f, k, G) = \text{cell}(G, i_2 -' 1, j_2)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\text{front_left_cell}(f, k, G) = \text{cell}(G, i_2, j_2)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\text{front_left_cell}(f, k, G) = \text{cell}(G, i_2 -' 1, j_2 -' 1)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\text{front_left_cell}(f, k, G) = \text{cell}(G, i_2, j_2 -' 1)$.

The following propositions are true:

- (35) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i, j+1 \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i, j+1)$. Then $\text{front_left_cell}(f, k, G) = \text{cell}(G, i -' 1, j+1)$.
- (36) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i, j+1 \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i, j+1)$. Then $\text{front_right_cell}(f, k, G) = \text{cell}(G, i, j+1)$.
- (37) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i+1, j \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i+1, j)$. Then $\text{front_left_cell}(f, k, G) = \text{cell}(G, i+1, j)$.
- (38) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i+1, j \rangle \in$ the indices of G and $f_k = G \circ (i, j)$ and $f_{k+1} = G \circ (i+1, j)$. Then $\text{front_right_cell}(f, k, G) = \text{cell}(G, i+1, j -' 1)$.
- (39) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i+1, j \rangle \in$ the indices of G and $f_k = G \circ (i+1, j)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{front_left_cell}(f, k, G) = \text{cell}(G, i -' 1, j -' 1)$.
- (40) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j \rangle \in$ the indices of G and $\langle i+1, j \rangle \in$ the indices of G and $f_k = G \circ (i+1, j)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{front_right_cell}(f, k, G) = \text{cell}(G, i -' 1, j)$.
- (41) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j+1 \rangle \in$ the indices of G and $\langle i, j \rangle \in$ the indices of G and $f_k = G \circ (i, j+1)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{front_left_cell}(f, k, G) = \text{cell}(G, i, j -' 1)$.
- (42) Suppose that $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $\langle i, j+1 \rangle \in$ the indices of G and $\langle i, j \rangle \in$ the indices of G and $f_k = G \circ (i, j+1)$ and $f_{k+1} = G \circ (i, j)$. Then $\text{front_right_cell}(f, k, G) = \text{cell}(G, i -' 1, j -' 1)$.
- (43) Suppose $1 \leq k$ and $k+1 \leq \text{len } f$ and f is a sequence which elements belong to G and $k+1 \leq n$. Then $\text{front_left_cell}(f, k, G) = \text{front_left_cell}(f \upharpoonright n, k, G)$ and $\text{front_right_cell}(f, k, G) = \text{front_right_cell}(f \upharpoonright n, k, G)$.

Let D be a set, let f be a finite sequence of elements of D , let G be a matrix over D , and let us consider k . We say that f turns right k , G if and only if the condition (Def. 6) is satisfied.

- (Def. 6) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\langle i_2 + 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 + 1, j_2)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\langle i_2, j_2 - 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 - 1)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\langle i_2, j_2 + 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 + 1)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\langle i_2 - 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 - 1, j_2)$.

We say that f turns left k , G if and only if the condition (Def. 7) is satisfied.

- (Def. 7) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\langle i_2 - 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 - 1, j_2)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\langle i_2, j_2 + 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 + 1)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\langle i_2, j_2 - 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 - 1)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\langle i_2 + 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 + 1, j_2)$.

We say that f goes straight k , G if and only if the condition (Def. 8) is satisfied.

- (Def. 8) Let i_1, j_1, i_2, j_2 be natural numbers. Suppose $\langle i_1, j_1 \rangle \in$ the indices of G and $\langle i_2, j_2 \rangle \in$ the indices of G and $f_k = G \circ (i_1, j_1)$ and $f_{k+1} = G \circ (i_2, j_2)$. Then
- (i) $i_1 = i_2$ and $j_1 + 1 = j_2$ and $\langle i_2, j_2 + 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 + 1)$, or
 - (ii) $i_1 + 1 = i_2$ and $j_1 = j_2$ and $\langle i_2 + 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 + 1, j_2)$, or
 - (iii) $i_1 = i_2 + 1$ and $j_1 = j_2$ and $\langle i_2 - 1, j_2 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2 - 1, j_2)$, or
 - (iv) $i_1 = i_2$ and $j_1 = j_2 + 1$ and $\langle i_2, j_2 - 1 \rangle \in$ the indices of G and $f_{k+2} = G \circ (i_2, j_2 - 1)$.

We adopt the following convention: D denotes a set, f, f_1, f_2 denote finite sequences of elements of D , and G denotes a matrix over D .

The following propositions are true:

- (44) If $1 \leq k$ and $k + 2 \leq \text{len } f$ and $k + 2 \leq n$ and $f \upharpoonright n$ turns right k , G , then f turns right k , G .
- (45) If $1 \leq k$ and $k + 2 \leq \text{len } f$ and $k + 2 \leq n$ and $f \upharpoonright n$ turns left k , G , then f turns left k , G .
- (46) If $1 \leq k$ and $k + 2 \leq \text{len } f$ and $k + 2 \leq n$ and $f \upharpoonright n$ goes straight k , G , then f goes straight k , G .
- (47) Suppose that $1 < k$ and $k + 1 \leq \text{len } f_1$ and $k + 1 \leq \text{len } f_2$ and f_1 is a sequence which elements belong to G and $f_1 \upharpoonright k = f_2 \upharpoonright k$ and f_1 turns right $k - 1$, G and f_2 turns right $k - 1$, G . Then $f_1 \upharpoonright (k + 1) = f_2 \upharpoonright (k + 1)$.
- (48) Suppose that $1 < k$ and $k + 1 \leq \text{len } f_1$ and $k + 1 \leq \text{len } f_2$ and f_1 is a sequence which elements belong to G and $f_1 \upharpoonright k = f_2 \upharpoonright k$ and f_1 turns left $k - 1$, G and f_2 turns left $k - 1$, G . Then $f_1 \upharpoonright (k + 1) = f_2 \upharpoonright (k + 1)$.
- (49) Suppose that $1 < k$ and $k + 1 \leq \text{len } f_1$ and $k + 1 \leq \text{len } f_2$ and f_1 is a sequence which elements belong to G and $f_1 \upharpoonright k = f_2 \upharpoonright k$ and f_1 goes straight $k - 1$, G and f_2 goes straight $k - 1$, G . Then $f_1 \upharpoonright (k + 1) = f_2 \upharpoonright (k + 1)$.
- (50) For every non empty set D and for every matrix M over D such that $1 \leq i$ and $i \leq \text{len } M$ and $1 \leq j$ and $j \leq \text{width } M$ holds $M \circ (i, j) \in \text{Values } M$.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/funct_6.html.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [8] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [9] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [10] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [11] Agata Darmochwał and Yatsuka Nakamura. The topological space \mathcal{E}_T^2 . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal1.html>.
- [12] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/matrix_1.html.
- [13] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/rfinseq.html>.
- [14] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard1.html>.
- [15] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part II. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard2.html>.
- [16] Yatsuka Nakamura and Andrzej Trybulec. Decomposing a Go-Board into cells. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard5.html>.
- [17] Andrzej Nędzusiak. σ -fields and probability. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/prob_1.html.
- [18] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/binarith.html>.
- [19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [20] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [21] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [22] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [23] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received April 23, 1999

Published January 2, 2004
