

Rotating and Reversing

Andrzej Trybulec
University of Białystok

Summary. Quite a number of lemmas for the Jordan curve theorem, as yet in the case of the special polygonal curves, have been proved. By “special” we mean, that it is a polygonal curve with edges parallel to axes and actually the lemmas have been proved, mostly, for the triangulations i.e. for finite sequences that define the curve. Moreover some of the results deal only with a special case:

- finite sequences are clockwise oriented,
- the first member of the sequence is the member with the lowest ordinate among those with the highest abscissa (N-min f , where f is a finite sequence, in the Mizar jargon).

In the change of the orientation one has to reverse the sequence (the operation introduced in [6]) and to change the second restriction one has to rotate the sequence (the operation introduced in [19]). The goal of the paper is to prove, mostly simple, facts about the relationship between properties and attributes of the finite sequence and its rotation (similar results about reversing had been proved in [6]). Some of them deal with recounting parameters, others with properties that are invariant under the rotation. We prove also that the finite sequence is either clockwise oriented or it is such after reversing. Everything is proved for the so called standard finite sequences, which means that if a point belongs to it then every point with the same abscissa or with the same ordinate, that belongs to the polygon, belongs also to the finite sequence. It does not seem that this requirement causes serious technical obstacles.

MML Identifier: REVROT_1.

WWW: http://mizar.org/JFM/Vol11/revrot_1.html

The articles [17], [22], [2], [15], [1], [4], [3], [5], [9], [21], [10], [6], [19], [16], [7], [8], [11], [12], [13], [18], [20], and [14] provide the notation and terminology for this paper.

1. PRELIMINARIES

For simplicity, we use the following convention: i, k, m, n are natural numbers, D is a non empty set, p is an element of D , and f is a finite sequence of elements of D .

Let S be a non trivial 1-sorted structure. One can verify that the carrier of S is non trivial.

Let D be a non empty set and let f be a finite sequence of elements of D . Let us observe that f is constant if and only if:

(Def. 1) For all n, m such that $n \in \text{dom } f$ and $m \in \text{dom } f$ holds $f_n = f_m$.

We now state three propositions:

- (1) Let D be a non empty set and f be a finite sequence of elements of D . If f yields $f_{\text{len } f}$ just once, then $f_{\text{len } f} \leftrightarrow^p f = \text{len } f$.
- (2) For every non empty set D and for every finite sequence f of elements of D holds $f_{|\text{len } f} = \emptyset$.

- (3) For every non empty set D and for every non empty finite sequence f of elements of D holds $f_{\text{len } f} \in \text{rng } f$.

Let D be a non empty set, let f be a finite sequence of elements of D , and let y be a set. Let us observe that f yields y just once if and only if:

- (Def. 2) There exists a set x such that $x \in \text{dom } f$ and $y = f_x$ and for every set z such that $z \in \text{dom } f$ and $z \neq x$ holds $f_z \neq y$.

One can prove the following propositions:

- (4) Let D be a non empty set and f be a finite sequence of elements of D . If f yields $f_{\text{len } f}$ just once, then $f^- : f_{\text{len } f} = f$.
- (5) Let D be a non empty set and f be a finite sequence of elements of D . If f yields $f_{\text{len } f}$ just once, then $f^- : f_{\text{len } f} = \langle f_{\text{len } f} \rangle$.
- (6) $1 \leq \text{len}(f^- p)$.
- (7) Let D be a non empty set, p be an element of D , and f be a finite sequence of elements of D . If $p \in \text{rng } f$, then $\text{len}(f^- p) \leq \text{len } f$.
- (8) For every non empty set D and for every circular non empty finite sequence f of elements of D holds $\text{Rev}(f)$ is circular.

2. ABOUT THE ROTATION

In the sequel D denotes a non empty set, p denotes an element of D , and f denotes a finite sequence of elements of D .

Next we state several propositions:

- (9) If $p \in \text{rng } f$ and $1 \leq i$ and $i \leq \text{len}(f^- p)$, then $(f \circ p)_i = f_{(i-1)+p \oplus f}$.
- (10) If $p \in \text{rng } f$ and $p \oplus f \leq i$ and $i \leq \text{len } f$, then $f_i = (f \circ p)_{(i+1)-p \oplus f}$.
- (11) If $p \in \text{rng } f$, then $(f \circ p)_{\text{len}(f^- p)} = f_{\text{len } f}$.
- (12) If $p \in \text{rng } f$ and $\text{len}(f^- p) < i$ and $i \leq \text{len } f$, then $(f \circ p)_i = f_{(i+p \oplus f)-\text{len } f}$.
- (13) If $p \in \text{rng } f$ and $1 < i$ and $i \leq p \oplus f$, then $f_i = (f \circ p)_{(i+\text{len } f)-p \oplus f}$.
- (14) $\text{len}(f \circ p) = \text{len } f$.
- (15) $\text{dom}(f \circ p) = \text{dom } f$.
- (16) Let D be a non empty set, f be a circular finite sequence of elements of D , and p be an element of D . If for every i such that $1 < i$ and $i < \text{len } f$ holds $f_i \neq f_1$, then $(f \circ p) \circ f_1 = f$.

3. ROTATING CIRCULAR ONES

In the sequel f denotes a circular finite sequence of elements of D .

Next we state two propositions:

- (17) If $p \in \text{rng } f$ and $\text{len}(f^- p) \leq i$ and $i \leq \text{len } f$, then $(f \circ p)_i = f_{(i+p \oplus f)-\text{len } f}$.
- (18) If $p \in \text{rng } f$ and $1 \leq i$ and $i \leq p \oplus f$, then $f_i = (f \circ p)_{(i+\text{len } f)-p \oplus f}$.

Let D be a non trivial set. Observe that there exists a finite sequence of elements of D which is non constant and circular.

Let D be a non trivial set, let p be an element of D , and let f be a non constant circular finite sequence of elements of D . One can check that $f \circ p$ is non constant.

4. FINITE SEQUENCE ON THE PLANE

The following proposition is true

(19) For every non empty natural number n holds $0_{\mathcal{E}_T^n} \neq 1_{\text{REAL } n}$.

Let n be a non empty natural number. One can verify that \mathcal{E}_T^n is non trivial.

In the sequel f, g denote finite sequences of elements of \mathcal{E}_T^2 .

The following four propositions are true:

(20) If $\text{rng } f \subseteq \text{rng } g$, then $\text{rng } \mathbf{X}\text{-coordinate}(f) \subseteq \text{rng } \mathbf{X}\text{-coordinate}(g)$.

(21) If $\text{rng } f = \text{rng } g$, then $\text{rng } \mathbf{X}\text{-coordinate}(f) = \text{rng } \mathbf{X}\text{-coordinate}(g)$.

(22) If $\text{rng } f \subseteq \text{rng } g$, then $\text{rng } \mathbf{Y}\text{-coordinate}(f) \subseteq \text{rng } \mathbf{Y}\text{-coordinate}(g)$.

(23) If $\text{rng } f = \text{rng } g$, then $\text{rng } \mathbf{Y}\text{-coordinate}(f) = \text{rng } \mathbf{Y}\text{-coordinate}(g)$.

5. ROTATING FINITE SEQUENCE ON THE PLANE

In the sequel p denotes a point of \mathcal{E}_T^2 and f denotes a finite sequence of elements of \mathcal{E}_T^2 .

Let p be a point of \mathcal{E}_T^2 and let f be a special circular finite sequence of elements of \mathcal{E}_T^2 . One can check that $f \circ p$ is special.

Next we state several propositions:

(24) If $p \in \text{rng } f$ and $1 \leq i$ and $i < \text{len}(f : -p)$, then $\mathcal{L}(f \circ p, i) = \mathcal{L}(f, (i - '1) + p \leftrightarrow f)$.

(25) If $p \in \text{rng } f$ and $p \leftrightarrow f \leq i$ and $i < \text{len } f$, then $\mathcal{L}(f, i) = \mathcal{L}(f \circ p, (i - 'p \leftrightarrow f) + 1)$.

(26) For every circular finite sequence f of elements of \mathcal{E}_T^2 holds $\text{Inc}(\mathbf{X}\text{-coordinate}(f)) = \text{Inc}(\mathbf{X}\text{-coordinate}(f \circ p))$.

(27) For every circular finite sequence f of elements of \mathcal{E}_T^2 holds $\text{Inc}(\mathbf{Y}\text{-coordinate}(f)) = \text{Inc}(\mathbf{Y}\text{-coordinate}(f \circ p))$.

(28) For every non empty circular finite sequence f of elements of \mathcal{E}_T^2 holds the Go-board of $f \circ p =$ the Go-board of f .

(29) For every non constant standard special circular sequence f holds $\text{Rev}(f \circ p) = \text{Rev}(f) \circ p$.

6. ROTATING CIRCULAR ONES (ON THE PLANE)

In the sequel f denotes a circular finite sequence of elements of \mathcal{E}_T^2 .

The following propositions are true:

(30) For every circular s.c.c. finite sequence f of elements of \mathcal{E}_T^2 such that $\text{len } f > 4$ holds $\mathcal{L}(f, \text{len } f - '1) \cap \mathcal{L}(f, 1) = \{f_1\}$.

(31) If $p \in \text{rng } f$ and $\text{len}(f : -p) \leq i$ and $i < \text{len } f$, then $\mathcal{L}(f \circ p, i) = \mathcal{L}(f, (i + p \leftrightarrow f) - ' \text{len } f)$.

Let p be a point of \mathcal{E}_T^2 and let f be a circular s.c.c. finite sequence of elements of \mathcal{E}_T^2 . One can check that $f \circ p$ is s.c.c..

Let p be a point of \mathcal{E}_T^2 and let f be a non constant standard special circular sequence. Observe that $f \circ p$ is unfolded.

One can prove the following propositions:

(32) If $p \in \text{rng } f$ and $1 \leq i$ and $i < p \leftrightarrow f$, then $\mathcal{L}(f, i) = \mathcal{L}(f \circ p, (i + \text{len } f) - ' p \leftrightarrow f)$.

(33) $\tilde{\mathcal{L}}(f \circ p) = \tilde{\mathcal{L}}(f)$.

- (34) Let G be a Go-board. Then f is a sequence which elements belong to G if and only if $f \circ p$ is a sequence which elements belong to G .

Let p be a point of \mathcal{E}_T^2 and let f be a standard non empty circular finite sequence of elements of \mathcal{E}_T^2 . Note that $f \circ p$ is standard.

Next we state three propositions:

- (35) Let f be a non constant standard special circular sequence and given p, k . If $p \in \text{rng } f$ and $1 \leq k$ and $k < p \leftarrow f$, then $\text{leftcell}(f, k) = \text{leftcell}(f \circ p, (k + \text{len } f) -' p \leftarrow f)$.
- (36) For every non constant standard special circular sequence f holds $\text{LeftComp}(f \circ p) = \text{LeftComp}(f)$.
- (37) For every non constant standard special circular sequence f holds $\text{RightComp}(f \circ p) = \text{RightComp}(f)$.

7. THE ORIENTATION

Let p be a point of \mathcal{E}_T^2 and let f be a clockwise oriented non constant standard special circular sequence. Observe that $f \circ p$ is clockwise oriented.

Next we state the proposition

- (38) Let f be a non constant standard special circular sequence. Then f is clockwise oriented or $\text{Rev}(f)$ is clockwise oriented.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [6] Czesław Byliński. Some properties of restrictions of finite sequences. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/finseq_5.html.
- [7] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [8] Agata Darmochwał and Yatsuka Nakamura. The topological space \mathcal{E}_T^2 . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal1.html>.
- [9] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/matrix_1.html.
- [10] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/rfinseq.html>.
- [11] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard1.html>.
- [12] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part II. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard2.html>.
- [13] Yatsuka Nakamura and Andrzej Trybulec. Decomposing a Go-Board into cells. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard5.html>.
- [14] Yatsuka Nakamura, Andrzej Trybulec, and Czesław Byliński. Bounded domains and unbounded domains. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/jordan2c.html>.
- [15] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/binarith.html>.
- [16] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.

- [17] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [18] Andrzej Trybulec. Left and right component of the complement of a special closed curve. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard9.html>.
- [19] Andrzej Trybulec. On the decomposition of finite sequences. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/finseq_6.html.
- [20] Andrzej Trybulec and Yatsuka Nakamura. On the order on a special polygon. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/sprect_2.html.
- [21] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [22] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received January 21, 1999

Published January 2, 2004
