

# Extremal Properties of Vertices on Special Polygons, Part I

Yatsuka Nakamura  
Shinshu University  
Nagano

Czesław Byliński  
Warsaw University  
Białystok

**Summary.** First, extremal properties of endpoints of line segments in  $n$ -dimensional Euclidean space are discussed. Some topological properties of line segments are also discussed. Secondly, extremal properties of vertices of special polygons which consist of horizontal and vertical line segments in 2-dimensional Euclidean space, are also derived.

MML Identifier: SPPOL\_1.

WWW: [http://mizar.org/JFM/Vol6/sppol\\_1.html](http://mizar.org/JFM/Vol6/sppol_1.html)

The articles [14], [18], [3], [15], [11], [2], [16], [6], [13], [8], [1], [4], [12], [5], [7], [17], [9], and [10] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

One can prove the following propositions:

- (2)<sup>1</sup> For every finite set  $A$  holds  $A$  is trivial iff  $\text{card}A < 2$ .
- (3) For every set  $A$  holds  $A$  is non trivial iff there exist sets  $a_1, a_2$  such that  $a_1 \in A$  and  $a_2 \in A$  and  $a_1 \neq a_2$ .
- (4) Let  $D$  be a set and  $A$  be a subset of  $D$ . Then  $A$  is non trivial if and only if there exist elements  $d_1, d_2$  of  $D$  such that  $d_1 \in A$  and  $d_2 \in A$  and  $d_1 \neq d_2$ .

We follow the rules:  $n, i, k, m$  denote natural numbers and  $r, r_1, r_2, s, s_1, s_2$  denote real numbers. One can prove the following propositions:

- (5) If  $r \leq s$ , then  $r - 1 \leq s$  and  $r - 1 < s$  and  $r \leq s + 1$  and  $r < s + 1$ .
- (6) If  $n < k$ , then  $n \leq k - 1$  and if  $r < s$ , then  $r - 1 \leq s$  and  $r - 1 < s$  and  $r \leq s + 1$  and  $r < s + 1$ .
- (7) If  $1 \leq k - m$  and  $k - m \leq n$ , then  $k - m \in \text{Seg}n$  and  $k - m$  is a natural number.
- (8) If  $r_1 \geq 0$  and  $r_2 \geq 0$  and  $r_1 + r_2 = 0$ , then  $r_1 = 0$  and  $r_2 = 0$ .
- (9) If  $r_1 \leq 0$  and  $r_2 \leq 0$  and  $r_1 + r_2 = 0$ , then  $r_1 = 0$  and  $r_2 = 0$ .
- (10) If  $0 \leq r_1$  and  $r_1 \leq 1$  and  $0 \leq r_2$  and  $r_2 \leq 1$  and  $r_1 \cdot r_2 = 1$ , then  $r_1 = 1$  and  $r_2 = 1$ .
- (11) If  $r_1 \geq 0$  and  $r_2 \geq 0$  and  $s_1 \geq 0$  and  $s_2 \geq 0$  and  $r_1 \cdot s_1 + r_2 \cdot s_2 = 0$ , then  $r_1 = 0$  or  $s_1 = 0$  but  $r_2 = 0$  or  $s_2 = 0$ .

---

<sup>1</sup> The proposition (1) has been removed.

- (12) If  $0 \leq r$  and  $r \leq 1$  and  $s_1 \geq 0$  and  $s_2 \geq 0$  and  $r \cdot s_1 + (1 - r) \cdot s_2 = 0$ , then  $r = 0$  and  $s_2 = 0$  or  $r = 1$  and  $s_1 = 0$  or  $s_1 = 0$  and  $s_2 = 0$ .
- (13) If  $r < r_1$  and  $r < r_2$ , then  $r < \min(r_1, r_2)$ .
- (14) If  $r > r_1$  and  $r > r_2$ , then  $r > \max(r_1, r_2)$ .

In this article we present several logical schemes. The scheme *FinSeqFam* deals with a non empty set  $\mathcal{A}$ , a finite sequence  $\mathcal{B}$  of elements of  $\mathcal{A}$ , a binary functor  $\mathcal{F}$  yielding a set, and a unary predicate  $\mathcal{P}$ , and states that:

$$\{\mathcal{F}(\mathcal{B}, i) : i \in \text{dom } \mathcal{B} \wedge \mathcal{P}[i]\} \text{ is finite}$$

for all values of the parameters.

The scheme *FinSeqFam'* deals with a non empty set  $\mathcal{A}$ , a finite sequence  $\mathcal{B}$  of elements of  $\mathcal{A}$ , a binary functor  $\mathcal{F}$  yielding a set, and a unary predicate  $\mathcal{P}$ , and states that:

$$\{\mathcal{F}(\mathcal{B}, i) : 1 \leq i \wedge i \leq \text{len } \mathcal{B} \wedge \mathcal{P}[i]\} \text{ is finite}$$

for all values of the parameters.

Next we state several propositions:

- (15) For all elements  $x_1, x_2, x_3$  of  $\mathcal{R}^n$  holds  $|x_1 - x_2| - |x_2 - x_3| \leq |x_1 - x_3|$ .
- (16) For all elements  $x_1, x_2, x_3$  of  $\mathcal{R}^n$  holds  $|x_2 - x_1| - |x_2 - x_3| \leq |x_3 - x_1|$ .
- (17) Every point of  $\mathcal{E}_T^n$  is an element of  $\mathcal{R}^n$  and a point of  $\mathcal{E}^n$ .
- (18) Every point of  $\mathcal{E}^n$  is an element of  $\mathcal{R}^n$  and a point of  $\mathcal{E}_T^n$ .
- (19) Every element of  $\mathcal{R}^n$  is a point of  $\mathcal{E}^n$  and a point of  $\mathcal{E}_T^n$ .

## 2. PROPERTIES OF LINE SEGMENTS

We use the following convention:  $r, s$  are real numbers and  $p, p_1, p_2, q_1, q_2$  are points of  $\mathcal{E}_T^n$ .

Next we state a number of propositions:

- (20) For all points  $u_1, u_2$  of  $\mathcal{E}^n$  and for all elements  $v_1, v_2$  of  $\mathcal{R}^n$  such that  $v_1 = u_1$  and  $v_2 = u_2$  holds  $\rho(u_1, u_2) = |v_1 - v_2|$ .
- (21) For all  $p, p_1, p_2$  such that  $p \in \mathcal{L}(p_1, p_2)$  there exists  $r$  such that  $0 \leq r$  and  $r \leq 1$  and  $p = (1 - r) \cdot p_1 + r \cdot p_2$ .
- (22) For all  $p_1, p_2, r$  such that  $0 \leq r$  and  $r \leq 1$  holds  $(1 - r) \cdot p_1 + r \cdot p_2 \in \mathcal{L}(p_1, p_2)$ .
- (23) Let  $P$  be a non empty subset of  $\mathcal{E}_T^n$ . Suppose  $P$  is closed and  $P \subseteq \mathcal{L}(p_1, p_2)$ . Then there exists  $s$  such that  $(1 - s) \cdot p_1 + s \cdot p_2 \in P$  and for every  $r$  such that  $0 \leq r$  and  $r \leq 1$  and  $(1 - r) \cdot p_1 + r \cdot p_2 \in P$  holds  $s \leq r$ .
- (24) For all  $p_1, p_2, q_1, q_2$  such that  $\mathcal{L}(q_1, q_2) \subseteq \mathcal{L}(p_1, p_2)$  and  $p_1 \in \mathcal{L}(q_1, q_2)$  holds  $p_1 = q_1$  or  $p_1 = q_2$ .
- (25) For all  $p_1, p_2, q_1, q_2$  such that  $\mathcal{L}(p_1, p_2) = \mathcal{L}(q_1, q_2)$  holds  $p_1 = q_1$  and  $p_2 = q_2$  or  $p_1 = q_2$  and  $p_2 = q_1$ .
- (26)  $\mathcal{E}_T^n$  is a  $T_2$  space.
- (27)  $\{p\}$  is closed.
- (28)  $\mathcal{L}(p_1, p_2)$  is compact.
- (29)  $\mathcal{L}(p_1, p_2)$  is closed.

Let us consider  $n, p$  and let  $P$  be a subset of  $\mathcal{E}_T^n$ . We say that  $p$  is extremal in  $P$  if and only if:

(Def. 1)  $p \in P$  and for all  $p_1, p_2$  such that  $p \in \mathcal{L}(p_1, p_2)$  and  $\mathcal{L}(p_1, p_2) \subseteq P$  holds  $p = p_1$  or  $p = p_2$ .

Next we state several propositions:

- (30) For all subsets  $P, Q$  of  $\mathcal{E}_T^n$  such that  $p$  is extremal in  $P$  and  $Q \subseteq P$  and  $p \in Q$  holds  $p$  is extremal in  $Q$ .
- (31)  $p$  is extremal in  $\{p\}$ .
- (32)  $p_1$  is extremal in  $\mathcal{L}(p_1, p_2)$ .
- (33)  $p_2$  is extremal in  $\mathcal{L}(p_1, p_2)$ .
- (34) If  $p$  is extremal in  $\mathcal{L}(p_1, p_2)$ , then  $p = p_1$  or  $p = p_2$ .

### 3. ALTERNATING SPECIAL SEQUENCES

We use the following convention:  $P, Q$  are subsets of  $\mathcal{E}_T^2$ ,  $f, f_1, f_2$  are finite sequences of elements of the carrier of  $\mathcal{E}_T^2$ , and  $p, p_1, p_2, p_3, q$  are points of  $\mathcal{E}_T^2$ .

One can prove the following proposition

- (35) For all  $p_1, p_2$  such that  $(p_1)_1 \neq (p_2)_1$  and  $(p_1)_2 \neq (p_2)_2$  there exists  $p$  such that  $p \in \mathcal{L}(p_1, p_2)$  and  $p_1 \neq (p)_1$  and  $p_1 \neq (p)_2$  and  $p_2 \neq (p)_1$  and  $p_2 \neq (p)_2$ .

Let us consider  $P$ . We say that  $P$  is horizontal if and only if:

- (Def. 2) For all  $p, q$  such that  $p \in P$  and  $q \in P$  holds  $p_2 = q_2$ .

We say that  $P$  is vertical if and only if:

- (Def. 3) For all  $p, q$  such that  $p \in P$  and  $q \in P$  holds  $p_1 = q_1$ .

Let us note that every subset of  $\mathcal{E}_T^2$  which is non trivial and horizontal is also non vertical and every subset of  $\mathcal{E}_T^2$  which is non trivial and vertical is also non horizontal.

One can prove the following propositions:

- (36)  $p_2 = q_2$  iff  $\mathcal{L}(p, q)$  is horizontal.
- (37)  $p_1 = q_1$  iff  $\mathcal{L}(p, q)$  is vertical.
- (38) If  $p_1 \in \mathcal{L}(p, q)$  and  $p_2 \in \mathcal{L}(p, q)$  and  $(p_1)_1 \neq (p_2)_1$  and  $(p_1)_2 = (p_2)_2$ , then  $\mathcal{L}(p, q)$  is horizontal.
- (39) If  $p_1 \in \mathcal{L}(p, q)$  and  $p_2 \in \mathcal{L}(p, q)$  and  $(p_1)_2 \neq (p_2)_2$  and  $(p_1)_1 = (p_2)_1$ , then  $\mathcal{L}(p, q)$  is vertical.
- (40)  $\mathcal{L}(f, i)$  is closed.
- (41) If  $f$  is special, then  $\mathcal{L}(f, i)$  is vertical or  $\mathcal{L}(f, i)$  is horizontal.
- (42) If  $f$  is one-to-one and  $1 \leq i$  and  $i+1 \leq \text{len } f$ , then  $\mathcal{L}(f, i)$  is non trivial.
- (43) If  $f$  is one-to-one and  $1 \leq i$  and  $i+1 \leq \text{len } f$  and  $\mathcal{L}(f, i)$  is vertical, then  $\mathcal{L}(f, i)$  is non horizontal.
- (44) For every  $f$  holds  $\{\mathcal{L}(f, i) : 1 \leq i \wedge i \leq \text{len } f\}$  is finite.
- (45) For every  $f$  holds  $\{\mathcal{L}(f, i) : 1 \leq i \wedge i+1 \leq \text{len } f\}$  is finite.
- (46) For every  $f$  holds  $\{\mathcal{L}(f, i) : 1 \leq i \wedge i \leq \text{len } f\}$  is a family of subsets of  $\mathcal{E}_T^2$ .
- (47) For every  $f$  holds  $\{\mathcal{L}(f, i) : 1 \leq i \wedge i+1 \leq \text{len } f\}$  is a family of subsets of  $\mathcal{E}_T^2$ .
- (48) For every  $f$  such that  $Q = \bigcup \{\mathcal{L}(f, i) : 1 \leq i \wedge i+1 \leq \text{len } f\}$  holds  $Q$  is closed.
- (49)  $\tilde{\mathcal{L}}(f)$  is closed.

Let  $I_1$  be a finite sequence of elements of  $\mathcal{E}_T^2$ . We say that  $I_1$  is alternating if and only if:

(Def. 4) For every  $i$  such that  $1 \leq i$  and  $i+2 \leq \text{len} I_1$  holds  $((I_1)_i)_1 \neq ((I_1)_{i+2})_1$  and  $((I_1)_i)_2 \neq ((I_1)_{i+2})_2$ .

The following propositions are true:

- (50) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $(f_i)_1 = (f_{i+1})_1$ , then  $(f_{i+1})_2 = (f_{i+2})_2$ .
- (51) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $(f_i)_2 = (f_{i+1})_2$ , then  $(f_{i+1})_1 = (f_{i+2})_1$ .
- (52) Suppose  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $p_1 = f_i$  and  $p_2 = f_{i+1}$  and  $p_3 = f_{i+2}$ . Then  $(p_1)_1 = (p_2)_1$  and  $(p_3)_1 \neq (p_2)_1$  or  $(p_1)_2 = (p_2)_2$  and  $(p_3)_2 \neq (p_2)_2$ .
- (53) Suppose  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $p_1 = f_i$  and  $p_2 = f_{i+1}$  and  $p_3 = f_{i+2}$ . Then  $(p_2)_1 = (p_3)_1$  and  $(p_1)_1 \neq (p_2)_1$  or  $(p_2)_2 = (p_3)_2$  and  $(p_1)_2 \neq (p_2)_2$ .
- (54) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$ , then  $\mathcal{L}(f_i, f_{i+2}) \not\subseteq \mathcal{L}(f, i) \cup \mathcal{L}(f, i+1)$ .
- (55) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $\mathcal{L}(f, i)$  is vertical, then  $\mathcal{L}(f, i+1)$  is horizontal.
- (56) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $\mathcal{L}(f, i)$  is horizontal, then  $\mathcal{L}(f, i+1)$  is vertical.
- (57) Suppose  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$ . Then  $\mathcal{L}(f, i)$  is vertical and  $\mathcal{L}(f, i+1)$  is horizontal or  $\mathcal{L}(f, i)$  is horizontal and  $\mathcal{L}(f, i+1)$  is vertical.
- (58) Suppose  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $f_{i+1} \in \mathcal{L}(p, q)$  and  $\mathcal{L}(p, q) \subseteq \mathcal{L}(f, i) \cup \mathcal{L}(f, i+1)$ . Then  $f_{i+1} = p$  or  $f_{i+1} = q$ .
- (59) If  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$ , then  $f_{i+1}$  is extremal in  $\mathcal{L}(f, i) \cup \mathcal{L}(f, i+1)$ .
- (60) Let  $u$  be a point of  $\mathcal{E}^2$ . Suppose  $f$  is special and alternating and  $1 \leq i$  and  $i+2 \leq \text{len} f$  and  $u = f_{i+1}$  and  $f_{i+1} \in \mathcal{L}(p, q)$  and  $f_{i+1} \neq q$  and  $p \notin \mathcal{L}(f, i) \cup \mathcal{L}(f, i+1)$ . Let given  $s$ . If  $s > 0$ , then there exists  $p_3$  such that  $p_3 \notin \mathcal{L}(f, i) \cup \mathcal{L}(f, i+1)$  and  $p_3 \in \mathcal{L}(p, q)$  and  $p_3 \in \text{Ball}(u, s)$ .

Let us consider  $f_1, f_2, P$ . We say that  $f_1$  and  $f_2$  are generators of  $P$  if and only if the conditions (Def. 5) are satisfied.

(Def. 5)  $f_1$  is alternating and special sequence and  $f_2$  is alternating and special sequence and  $(f_1)_1 = (f_2)_1$  and  $(f_1)_{\text{len} f_1} = (f_2)_{\text{len} f_2}$  and  $\langle (f_1)_2, (f_1)_1, (f_2)_2 \rangle$  is alternating and  $\langle (f_1)_{\text{len} f_1 - 1}, (f_1)_{\text{len} f_1}, (f_2)_{\text{len} f_2 - 1} \rangle$  is alternating and  $(f_1)_1 \neq (f_1)_{\text{len} f_1}$  and  $\tilde{\mathcal{L}}(f_1) \cap \tilde{\mathcal{L}}(f_2) = \{(f_1)_1, (f_1)_{\text{len} f_1}\}$  and  $P = \tilde{\mathcal{L}}(f_1) \cup \tilde{\mathcal{L}}(f_2)$ .

We now state the proposition

- (61) If  $f_1$  and  $f_2$  are generators of  $P$  and  $1 < i$  and  $i < \text{len} f_1$ , then  $(f_1)_i$  is extremal in  $P$ .

## REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll1/card\\_1.html](http://mizar.org/JFM/Voll1/card_1.html).
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll1/nat\\_1.html](http://mizar.org/JFM/Voll1/nat_1.html).
- [3] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll1/ordinal1.html>.

- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finseq\\_1.html](http://mizar.org/JFM/Vol1/finseq_1.html).
- [5] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/realset1.html>.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [7] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/compts\\_1.html](http://mizar.org/JFM/Vol1/compts_1.html).
- [8] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finset\\_1.html](http://mizar.org/JFM/Vol1/finset_1.html).
- [9] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [10] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal1.html>.
- [11] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/real\\_1.html](http://mizar.org/JFM/Vol1/real_1.html).
- [12] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/metric\\_1.html](http://mizar.org/JFM/Vol2/metric_1.html).
- [13] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/pre\\_topc.html](http://mizar.org/JFM/Vol1/pre_topc.html).
- [14] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [15] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [16] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/square\\_1.html](http://mizar.org/JFM/Vol1/square_1.html).
- [17] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/finseq\\_4.html](http://mizar.org/JFM/Vol2/finseq_4.html).
- [18] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).

*Received May 11, 1994*

*Published January 2, 2004*

---