

# Closure Operators and Subalgebras<sup>1</sup>

Grzegorz Bancerek  
Warsaw University  
Białystok

MML Identifier: WAYBEL10.

WWW: <http://mizar.org/JFM/Vol9/waybel10.html>

The articles [13], [8], [15], [16], [17], [5], [7], [6], [1], [12], [14], [2], [10], [3], [4], [11], [18], and [9] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

In this article we present several logical schemes. The scheme *SubrelstrEx* deals with a non empty relational structure  $\mathcal{A}$ , a set  $\mathcal{B}$ , and a unary predicate  $\mathcal{P}$ , and states that:

There exists a non empty full strict relational substructure  $S$  of  $\mathcal{A}$  such that for every element  $x$  of  $\mathcal{A}$  holds  $x$  is an element of  $S$  if and only if  $\mathcal{P}[x]$

provided the following conditions are met:

- $\mathcal{P}[\mathcal{B}]$ , and
- $\mathcal{B} \in$  the carrier of  $\mathcal{A}$ .

The scheme *RelstrEq* deals with non empty relational structures  $\mathcal{A}$ ,  $\mathcal{B}$ , a unary predicate  $\mathcal{P}$ , and a binary predicate  $Q$ , and states that:

The relational structure of  $\mathcal{A} =$  the relational structure of  $\mathcal{B}$

provided the following conditions are met:

- For every set  $x$  holds  $x$  is an element of  $\mathcal{A}$  iff  $\mathcal{P}[x]$ ,
- For every set  $x$  holds  $x$  is an element of  $\mathcal{B}$  iff  $\mathcal{P}[x]$ ,
- For all elements  $a, b$  of  $\mathcal{A}$  holds  $a \leq b$  iff  $Q[a, b]$ , and
- For all elements  $a, b$  of  $\mathcal{B}$  holds  $a \leq b$  iff  $Q[a, b]$ .

The scheme *SubrelstrEq1* deals with a non empty relational structure  $\mathcal{A}$ , non empty full relational substructures  $\mathcal{B}$ ,  $\mathcal{C}$  of  $\mathcal{A}$ , and a unary predicate  $\mathcal{P}$ , and states that:

The relational structure of  $\mathcal{B} =$  the relational structure of  $\mathcal{C}$

provided the following conditions are met:

- For every set  $x$  holds  $x$  is an element of  $\mathcal{B}$  iff  $\mathcal{P}[x]$ , and
- For every set  $x$  holds  $x$  is an element of  $\mathcal{C}$  iff  $\mathcal{P}[x]$ .

The scheme *SubrelstrEq2* deals with a non empty relational structure  $\mathcal{A}$ , non empty full relational substructures  $\mathcal{B}$ ,  $\mathcal{C}$  of  $\mathcal{A}$ , and a unary predicate  $\mathcal{P}$ , and states that:

The relational structure of  $\mathcal{B} =$  the relational structure of  $\mathcal{C}$

provided the parameters have the following properties:

- For every element  $x$  of  $\mathcal{A}$  holds  $x$  is an element of  $\mathcal{B}$  iff  $\mathcal{P}[x]$ , and
- For every element  $x$  of  $\mathcal{A}$  holds  $x$  is an element of  $\mathcal{C}$  iff  $\mathcal{P}[x]$ .

Next we state three propositions:

---

<sup>1</sup>This work has been partially supported by the Office of Naval Research Grant N00014-95-1-1336.

(1) For all binary relations  $R, Q$  holds  $R \subseteq Q$  iff  $R^\sim \subseteq Q^\sim$  and  $R^\sim \subseteq Q$  iff  $R \subseteq Q^\sim$ .

(3)<sup>1</sup> Let  $L, S$  be relational structures. Then

- (i)  $S$  is a relational substructure of  $L$  iff  $S^{\text{op}}$  is a relational substructure of  $L^{\text{op}}$ , and
- (ii)  $S^{\text{op}}$  is a relational substructure of  $L$  iff  $S$  is a relational substructure of  $L^{\text{op}}$ .

(4) Let  $L, S$  be relational structures. Then

- (i)  $S$  is a full relational substructure of  $L$  iff  $S^{\text{op}}$  is a full relational substructure of  $L^{\text{op}}$ , and
- (ii)  $S^{\text{op}}$  is a full relational substructure of  $L$  iff  $S$  is a full relational substructure of  $L^{\text{op}}$ .

Let  $L$  be a relational structure and let  $S$  be a full relational substructure of  $L$ . Then  $S^{\text{op}}$  is a strict full relational substructure of  $L^{\text{op}}$ .

Let  $X$  be a set and let  $L$  be a non empty relational structure. Observe that  $X \mapsto L$  is nonempty.

Let  $S$  be a relational structure and let  $T$  be a non empty reflexive relational structure. One can verify that there exists a map from  $S$  into  $T$  which is monotone.

Let  $L$  be a non empty relational structure. Note that every map from  $L$  into  $L$  which is projection is also monotone and idempotent.

Let  $S, T$  be non empty reflexive relational structures and let  $f$  be a monotone map from  $S$  into  $T$ . Note that  $f^\circ$  is monotone.

Let  $L$  be a 1-sorted structure. Observe that  $\text{id}_L$  is one-to-one.

Let  $L$  be a non empty reflexive relational structure. Observe that  $\text{id}_L$  is sups-preserving and infs-preserving.

We now state the proposition

(5) Let  $L$  be a relational structure and  $S$  be a subset of  $L$ . Then  $\text{id}_S$  is a map from  $\text{sub}(S)$  into  $L$  and for every map  $f$  from  $\text{sub}(S)$  into  $L$  such that  $f = \text{id}_S$  holds  $f$  is monotone.

Let  $L$  be a non empty reflexive relational structure. One can verify that there exists a map from  $L$  into  $L$  which is sups-preserving, infs-preserving, closure, kernel, and one-to-one.

One can prove the following proposition

(6) Let  $L$  be a non empty reflexive relational structure,  $c$  be a closure map from  $L$  into  $L$ , and  $x$  be an element of  $L$ . Then  $c(x) \geq x$ .

Let  $S, T$  be 1-sorted structures, let  $f$  be a function from the carrier of  $S$  into the carrier of  $T$ , and let  $R$  be a 1-sorted structure. Let us assume that the carrier of  $R \subseteq$  the carrier of  $S$ . The functor  $f \upharpoonright R$  yielding a map from  $R$  into  $T$  is defined by:

(Def. 1)  $f \upharpoonright R = f \upharpoonright \text{the carrier of } R$ .

The following two propositions are true:

(7) Let  $S, T$  be relational structures,  $R$  be a relational substructure of  $S$ , and  $f$  be a function from the carrier of  $S$  into the carrier of  $T$ . Then  $f \upharpoonright R = f \upharpoonright \text{the carrier of } R$  and for every set  $x$  such that  $x \in$  the carrier of  $R$  holds  $(f \upharpoonright R)(x) = f(x)$ .

(8) Let  $S, T$  be relational structures and  $f$  be a map from  $S$  into  $T$ . Suppose  $f$  is one-to-one. Let  $R$  be a relational substructure of  $S$ . Then  $f \upharpoonright R$  is one-to-one.

Let  $S, T$  be non empty reflexive relational structures, let  $f$  be a monotone map from  $S$  into  $T$ , and let  $R$  be a relational substructure of  $S$ . One can verify that  $f \upharpoonright R$  is monotone.

Next we state the proposition

(9) Let  $S, T$  be non empty relational structures,  $R$  be a non empty relational substructure of  $S$ ,  $f$  be a map from  $S$  into  $T$ , and  $g$  be a map from  $T$  into  $S$ . Suppose  $f$  is one-to-one and  $g = f^{-1}$ . Then  $g \upharpoonright \text{Im}(f \upharpoonright R)$  is a map from  $\text{Im}(f \upharpoonright R)$  into  $R$  and  $g \upharpoonright \text{Im}(f \upharpoonright R) = (f \upharpoonright R)^{-1}$ .

<sup>1</sup> The proposition (2) has been removed.

## 2. THE LATTICE OF CLOSURE OPERATORS

Let  $S$  be a relational structure and let  $T$  be a non empty reflexive relational structure. Observe that  $\text{MonMaps}(S, T)$  is non empty.

One can prove the following proposition

- (10) Let  $S$  be a relational structure,  $T$  be a non empty reflexive relational structure, and  $x$  be a set. Then  $x$  is an element of  $\text{MonMaps}(S, T)$  if and only if  $x$  is a monotone map from  $S$  into  $T$ .

Let  $L$  be a non empty reflexive relational structure. The functor  $\text{CLOpers}(L)$  yields a non empty full strict relational substructure of  $\text{MonMaps}(L, L)$  and is defined by:

(Def. 2) For every map  $f$  from  $L$  into  $L$  holds  $f$  is an element of  $\text{CLOpers}(L)$  iff  $f$  is closure.

The following propositions are true:

- (11) Let  $L$  be a non empty reflexive relational structure and  $x$  be a set. Then  $x$  is an element of  $\text{CLOpers}(L)$  if and only if  $x$  is a closure map from  $L$  into  $L$ .
- (12) Let  $X$  be a set,  $L$  be a non empty relational structure,  $f, g$  be functions from  $X$  into the carrier of  $L$ , and  $x, y$  be elements of  $L^X$ . If  $x = f$  and  $y = g$ , then  $x \leq y$  iff  $f \leq g$ .
- (13) Let  $L$  be a complete lattice,  $c_1, c_2$  be maps from  $L$  into  $L$ , and  $x, y$  be elements of  $\text{CLOpers}(L)$ . If  $x = c_1$  and  $y = c_2$ , then  $x \leq y$  iff  $c_1 \leq c_2$ .
- (14) Let  $L$  be a reflexive relational structure and  $S_1, S_2$  be full relational substructures of  $L$ . Suppose the carrier of  $S_1 \subseteq$  the carrier of  $S_2$ . Then  $S_1$  is a relational substructure of  $S_2$ .
- (15) Let  $L$  be a complete lattice and  $c_1, c_2$  be closure maps from  $L$  into  $L$ . Then  $c_1 \leq c_2$  if and only if  $\text{Im } c_2$  is a relational substructure of  $\text{Im } c_1$ .

## 3. THE LATTICE OF CLOSURE SYSTEMS

Let  $L$  be a relational structure. The functor  $\text{Sub}(L)$  yields a strict non empty relational structure and is defined by the conditions (Def. 3).

- (Def. 3)(i) For every set  $x$  holds  $x$  is an element of  $\text{Sub}(L)$  iff  $x$  is a strict relational substructure of  $L$ , and
- (ii) for all elements  $a, b$  of  $\text{Sub}(L)$  holds  $a \leq b$  iff there exists a relational structure  $R$  such that  $b = R$  and  $a$  is a relational substructure of  $R$ .

Next we state the proposition

- (16) Let  $L, R$  be relational structures and  $x, y$  be elements of  $\text{Sub}(L)$ . Suppose  $y = R$ . Then  $x \leq y$  if and only if  $x$  is a relational substructure of  $R$ .

Let  $L$  be a relational structure. Note that  $\text{Sub}(L)$  is reflexive, antisymmetric, and transitive.

Let  $L$  be a relational structure. One can verify that  $\text{Sub}(L)$  is complete.

Let  $L$  be a complete lattice. One can verify that every relational substructure of  $L$  which is infs-inheriting is also non empty and every relational substructure of  $L$  which is sups-inheriting is also non empty.

Let  $L$  be a relational structure. A system of  $L$  is a full relational substructure of  $L$ .

Let  $L$  be a non empty relational structure and let  $S$  be a system of  $L$ . We introduce  $S$  is closure as a synonym of  $S$  is infs-inheriting.

Let  $L$  be a non empty relational structure. Observe that  $\text{sub}(\Omega_L)$  is infs-inheriting and sups-inheriting.

Let  $L$  be a non empty relational structure. The functor  $\text{ClosureSystems}(L)$  yields a full strict non empty relational substructure of  $\text{Sub}(L)$  and is defined by the condition (Def. 4).

(Def. 4) Let  $R$  be a strict relational substructure of  $L$ . Then  $R$  is an element of  $\text{ClosureSystems}(L)$  if and only if  $R$  is infs-inheriting and full.

One can prove the following two propositions:

- (17) Let  $L$  be a non empty relational structure and  $x$  be a set. Then  $x$  is an element of  $\text{ClosureSystems}(L)$  if and only if  $x$  is a strict closure system of  $L$ .
- (18) Let  $L$  be a non empty relational structure,  $R$  be a relational structure, and  $x, y$  be elements of  $\text{ClosureSystems}(L)$ . Suppose  $y = R$ . Then  $x \leq y$  if and only if  $x$  is a relational substructure of  $R$ .

#### 4. ISOMORPHISM BETWEEN CLOSURE OPERATORS AND CLOSURE SYSTEMS

Let  $L$  be a non empty poset and let  $h$  be a closure map from  $L$  into  $L$ . One can verify that  $\text{Im } h$  is infs-inheriting.

Let  $L$  be a non empty poset. The functor  $\text{CIImageMap}(L)$  yielding a map from  $\text{CIOPers}(L)$  into  $(\text{ClosureSystems}(L))^{\text{op}}$  is defined as follows:

(Def. 5) For every closure map  $c$  from  $L$  into  $L$  holds  $(\text{CIImageMap}(L))(c) = \text{Im } c$ .

Let  $L$  be a non empty relational structure and let  $S$  be a relational substructure of  $L$ . The closure operation of  $S$  is a map from  $L$  into  $L$  and is defined by:

(Def. 6) For every element  $x$  of  $L$  holds (the closure operation of  $S$ )( $x$ ) =  $\bigcap_L(\uparrow x \cap \text{the carrier of } S)$ .

Let  $L$  be a complete lattice and let  $S$  be a closure system of  $L$ . Note that the closure operation of  $S$  is closure.

Next we state two propositions:

- (19) Let  $L$  be a complete lattice and  $S$  be a closure system of  $L$ . Then  $\text{Im}(\text{the closure operation of } S) = \text{the relational structure of } S$ .
- (20) For every complete lattice  $L$  and for every closure map  $c$  from  $L$  into  $L$  holds the closure operation of  $\text{Im } c = c$ .

Let  $L$  be a complete lattice. Note that  $\text{CIImageMap}(L)$  is one-to-one.

Next we state two propositions:

- (21) For every complete lattice  $L$  holds  $(\text{CIImageMap}(L))^{-1}$  is a map from  $(\text{ClosureSystems}(L))^{\text{op}}$  into  $\text{CIOPers}(L)$ .
- (22) Let  $L$  be a complete lattice and  $S$  be a strict closure system of  $L$ . Then  $(\text{CIImageMap}(L))^{-1}(S) = \text{the closure operation of } S$ .

Let  $L$  be a complete lattice. Note that  $\text{CIImageMap}(L)$  is isomorphic.

One can prove the following proposition

- (23) For every complete lattice  $L$  holds  $\text{CIOPers}(L)$  and  $(\text{ClosureSystems}(L))^{\text{op}}$  are isomorphic.

#### 5. ISOMORPHISM BETWEEN CLOSURE OPERATORS PRESERVING DIRECTED SUPS AND SUBALGEBRAS

One can prove the following three propositions:

- (24) Let  $L$  be a relational structure,  $S$  be a full relational substructure of  $L$ , and  $X$  be a subset of  $S$ . Then
- (i) if  $X$  is a directed subset of  $L$ , then  $X$  is directed, and
  - (ii) if  $X$  is a filtered subset of  $L$ , then  $X$  is filtered.

- (25) Let  $L$  be a complete lattice and  $S$  be a closure system of  $L$ . Then the closure operation of  $S$  is directed-sups-preserving if and only if  $S$  is directed-sups-inheriting.
- (26) Let  $L$  be a complete lattice and  $h$  be a closure map from  $L$  into  $L$ . Then  $h$  is directed-sups-preserving if and only if  $\text{Im } h$  is directed-sups-inheriting.

Let  $L$  be a complete lattice and let  $S$  be a directed-sups-inheriting closure system of  $L$ . Note that the closure operation of  $S$  is directed-sups-preserving.

Let  $L$  be a complete lattice and let  $h$  be a directed-sups-preserving closure map from  $L$  into  $L$ . Observe that  $\text{Im } h$  is directed-sups-inheriting.

Let  $L$  be a non empty reflexive relational structure. The functor  $\text{ClOps}^*(L)$  yields a non empty full strict relational substructure of  $\text{ClOps}(L)$  and is defined by the condition (Def. 7).

- (Def. 7) Let  $f$  be a closure map from  $L$  into  $L$ . Then  $f$  is an element of  $\text{ClOps}^*(L)$  if and only if  $f$  is directed-sups-preserving.

We now state the proposition

- (27) Let  $L$  be a non empty reflexive relational structure and  $x$  be a set. Then  $x$  is an element of  $\text{ClOps}^*(L)$  if and only if  $x$  is a directed-sups-preserving closure map from  $L$  into  $L$ .

Let  $L$  be a non empty relational structure. The functor  $\text{Subalgebras}(L)$  yields a full strict non empty relational substructure of  $\text{ClosureSystems}(L)$  and is defined by the condition (Def. 8).

- (Def. 8) Let  $R$  be a strict closure system of  $L$ . Then  $R$  is an element of  $\text{Subalgebras}(L)$  if and only if  $R$  is directed-sups-inheriting.

The following two propositions are true:

- (28) Let  $L$  be a non empty relational structure and  $x$  be a set. Then  $x$  is an element of  $\text{Subalgebras}(L)$  if and only if  $x$  is a strict directed-sups-inheriting closure system of  $L$ .
- (29) For every complete lattice  $L$  holds  $\text{Im}(\text{ClImageMap}(L) \upharpoonright \text{ClOps}^*(L)) = (\text{Subalgebras}(L))^{\text{op}}$ .

Let  $L$  be a complete lattice. Note that  $(\text{ClImageMap}(L) \upharpoonright \text{ClOps}^*(L))^{\circ}$  is isomorphic.

The following proposition is true

- (30) For every complete lattice  $L$  holds  $\text{ClOps}^*(L)$  and  $(\text{Subalgebras}(L))^{\text{op}}$  are isomorphic.

## REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/card\\_3.html](http://mizar.org/JFM/Vol2/card_3.html).
- [2] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [3] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_0.html](http://mizar.org/JFM/Vol8/yellow_0.html).
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_0.html](http://mizar.org/JFM/Vol8/waybel_0.html).
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [7] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [8] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [9] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_1.html](http://mizar.org/JFM/Vol8/waybel_1.html).
- [10] Adam Grabowski. On the category of posets. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/orders\\_3.html](http://mizar.org/JFM/Vol8/orders_3.html).
- [11] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_1.html](http://mizar.org/JFM/Vol8/yellow_1.html).

- [12] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. [http://mizar.org/JFM/Vol15/pralg\\_1.html](http://mizar.org/JFM/Vol15/pralg_1.html).
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [14] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/orders\\_1.html](http://mizar.org/JFM/Vol1/orders_1.html).
- [15] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [16] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).
- [17] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relset\\_1.html](http://mizar.org/JFM/Vol1/relset_1.html).
- [18] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_2.html](http://mizar.org/JFM/Vol8/yellow_2.html).

*Received January 15, 1997*

*Published January 2, 2004*

---