JOURNAL OF FORMALIZED MATHEMATICS
Volume Addenda,Released 1998,Published 2003
Inst. of Computer Science, Univ. of Białystok

Non-Negative Real Numbers. Part II[1]

Andrzej Trybulec
University of Białystok

MML Identifier: ARYTM_1.

WWW: http://mizar.org/JFM/Addenda/arytm_1.html

The articles [2], [4], [1], and [3] provide the notation and terminology for this paper.

In this paper x, y, z denote elements of $\mathbb{R}_+$.

Next we state several propositions:

(1) If $x+y=y$, then $x=\emptyset$.

(2) If $x*y=\emptyset$, then $x=\emptyset$ or $y=\emptyset$.

(3) If $x\leq y$ and $y\leq z$, then $x\leq z$.

(4) If $x\leq y$ and $y\leq x$, then $x=y$.

(5) If $x\leq y$ and $y=\emptyset$, then $x=\emptyset$.

(6) If $x=\emptyset$, then $x\leq y$.

(7) $x\leq y$ iff $x+z\leq y+z$.

(8) If $x\leq y$, then $x*z\leq y*z$.

Let x, y be elements of $\mathbb{R}_+$. The functor $x-'y$ yields an element of $\mathbb{R}_+$ and is defined as follows:

(Def. 1)(i) $(x-'y)+y=x$ if $y\leq x$,

(ii) $x-'y=\emptyset$, otherwise.

The following propositions are true:

(9) $x\leq y$ or $x-'y\neq\emptyset$.

(10) If $x\leq y$ and $y-'x=\emptyset$, then $x=y$.

(11) $x-'y\leq x$.

(12) If $y\leq x$ and $y\leq z$, then $x+(z-'y)=(x-'y)+z$.

(13) If $z\leq y$, then $x+(y-'z)=(x+y)-'z$.

(14) If $z\leq x$ and $y\leq z$, then $(x-'z)+y=x-'(z-'y)$.

(15) If $y\leq x$ and $y\leq z$, then $(z-'y)+x=(x-'y)+z$.

[1]This work has been supported by KBN Grant 8 T11C 018 12.

© Association of Mizar Users

(16) If $x \leq y$, then $z -' y \leq z -' x$.

(17) If $x \leq y$, then $x -' z \leq y -' z$.

Let x, y be elements of $\mathbb{R}_+$. The functor $x - y$ is defined as follows:

(Def. 2) $x - y = \begin{cases} \text{(i)} & x -' y, \text{ if } y \leq x, \\ & \langle \emptyset, y -' x \rangle, \text{ otherwise.} \end{cases}$

The following propositions are true:

(18) $x - x = \emptyset$.

(19) If $x = \emptyset$ and $y \neq \emptyset$, then $x - y = \langle \emptyset, y \rangle$.

(20) If $z \leq y$, then $x + (y -' z) = (x + y) - z$.

(21) If $z \not\leq y$, then $x - (z -' y) = (x + y) - z$.

(22) If $y \leq x$ and $y \not\leq z$, then $x - (y -' z) = (x -' y) + z$.

(23) If $y \not\leq x$ and $y \not\leq z$, then $x - (y -' z) = z - (y -' x)$.

(24) If $y \leq x$, then $x - (y + z) = (x -' y) - z$.

(25) If $x \leq y$ and $z \leq y$, then $(y -' z) - x = (y -' x) - z$.

(26) If $z \leq y$, then $x * (y -' z) = x * y - x * z$.

(27) If $z \not\leq y$ and $x \neq \emptyset$, then $\langle \emptyset, x * (z -' y) \rangle = x * y - x * z$.

(28) If $y -' z \neq \emptyset$ and $z \leq y$ and $x \neq \emptyset$, then $x * z - x * y = \langle \emptyset, x * (y -' z) \rangle$.

REFERENCES

[1] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal2.html.

[2] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[3] Andrzej Trybulec. Non negative real numbers. Part I. *Journal of Formalized Mathematics*, Addenda, 1998. http://mizar.org/JFM/Addenda/arytm_2.html.

[4] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received March 7, 1998

Published January 2, 2004