JOURNAL OF FORMALIZED MATHEMATICS
Volume 11, Released 1999, Published 2003
Inst. of Computer Science, Univ. of Białystok

Predicate Calculus for Boolean Valued Functions. Part II

Shunichi Kobayashi
Shinshu University
Nagano

Yatsuka Nakamura
Shinshu University
Nagano

Summary. In this paper, we have proved some elementary predicate calculus formulae containing the quantifiers of Boolean valued functions with respect to partitions. Such a theory is an analogy of usual predicate logic.

MML Identifier: BVFUNC_4.

WWW: http://mizar.org/JFM/Vol11/bvfunc_4.html

The articles [5], [7], [10], [9], [8], [1], [6], [4], [2], and [3] provide the notation and terminology for this paper.

1. Preliminaries

In this paper Y is a non empty set.

The following propositions are true:

(1) For all elements a, b, c of $\mathit{Boolean}^Y$ such that $a \Subset b \Rightarrow c$ holds $a \wedge b \Subset c$.

(2) For all elements a, b, c of $\mathit{Boolean}^Y$ such that $a \wedge b \Subset c$ holds $a \Subset b \Rightarrow c$.

(3) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \vee a \wedge b = a$.

(4) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \wedge (a \vee b) = a$.

(5) For every element a of $\mathit{Boolean}^Y$ holds $a \wedge \neg a = \mathit{false}(Y)$.

(6) For every element a of $\mathit{Boolean}^Y$ holds $a \vee \neg a = \mathit{true}(Y)$.

(7) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \Leftrightarrow b = (a \Rightarrow b) \wedge (b \Rightarrow a)$.

(8) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \Rightarrow b = \neg a \vee b$.

(9) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \oplus b = \neg a \wedge b \vee a \wedge \neg b$.

(10) For all elements a, b of $\mathit{Boolean}^Y$ holds $a \Leftrightarrow b = \mathit{true}(Y)$ iff $a \Rightarrow b = \mathit{true}(Y)$ and $b \Rightarrow a = \mathit{true}(Y)$.

(11) For all elements a, b, c of $\mathit{Boolean}^Y$ such that $a \Leftrightarrow b = \mathit{true}(Y)$ and $b \Leftrightarrow c = \mathit{true}(Y)$ holds $a \Leftrightarrow c = \mathit{true}(Y)$.

(12) For all elements a, b of $\mathit{Boolean}^Y$ such that $a \Leftrightarrow b = \mathit{true}(Y)$ holds $\neg a \Leftrightarrow \neg b = \mathit{true}(Y)$.

(13) For all elements a, b, c, d of $Boolean^Y$ such that $a \Leftrightarrow b = true(Y)$ and $c \Leftrightarrow d = true(Y)$ holds $a \wedge c \Leftrightarrow b \wedge d = true(Y)$.

(14) For all elements a, b, c, d of $Boolean^Y$ such that $a \Leftrightarrow b = true(Y)$ and $c \Leftrightarrow d = true(Y)$ holds $a \Rightarrow c \Leftrightarrow b \Rightarrow d = true(Y)$.

(15) For all elements a, b, c, d of $Boolean^Y$ such that $a \Leftrightarrow b = true(Y)$ and $c \Leftrightarrow d = true(Y)$ holds $a \vee c \Leftrightarrow b \vee d = true(Y)$.

(16) For all elements a, b, c, d of $Boolean^Y$ such that $a \Leftrightarrow b = true(Y)$ and $c \Leftrightarrow d = true(Y)$ holds $a \Leftrightarrow c \Leftrightarrow b \Leftrightarrow d = true(Y)$.

2. PREDICATE CALCULUS

We now state a number of propositions:

(17) Let a, b be elements of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. Then $\forall_{a \Leftrightarrow b, P_1} G = \forall_{a \Rightarrow b, P_1} G \wedge \forall_{b \Rightarrow a, P_1} G$.

(18) For every element a of $Boolean^Y$ and for every subset G of PARTITIONS(Y) and for all partitions P_1, P_2 of Y holds $\forall_{a,P_1} G \Subset \exists_{a,P_2} G$.

(19) Let a, u be elements of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. Suppose G is independent and $P_1 \in G$ and u is independent of P_1, G. If $a \Rightarrow u = true(Y)$, then $\forall_{a,P_1} G \Rightarrow u = true(Y)$.

(20) Let u be an element of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. If u is independent of P_1, G, then $\exists_{u,P_1} G \Subset u$.

(21) Let u be an element of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. If u is independent of P_1, G, then $u \Subset \forall_{u,P_1} G$.

(22) Let u be an element of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1, P_2 be partitions of Y. If u is independent of P_2, G, then $\forall_{u,P_1} G \Subset \forall_{u,P_2} G$.

(23) Let u be an element of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1, P_2 be partitions of Y. If u is independent of P_1, G, then $\exists_{u,P_1} G \Subset \exists_{u,P_2} G$.

(24) For all elements a, b of $Boolean^Y$ and for every subset G of PARTITIONS(Y) and for every partition P_1 of Y holds $\forall_{a \Leftrightarrow b, P_1} G \Subset \forall_{a,P_1} G \Leftrightarrow \forall_{b,P_1} G$.

(25) For all elements a, b of $Boolean^Y$ and for every subset G of PARTITIONS(Y) and for every partition P_1 of Y holds $\forall_{a \wedge b, P_1} G \Subset a \wedge \forall_{b,P_1} G$.

(26) For all elements a, u of $Boolean^Y$ and for every subset G of PARTITIONS(Y) and for every partition P_1 of Y holds $\forall_{a,P_1} G \Rightarrow u \Subset \exists_{a \Rightarrow u, P_1} G$.

(27) Let a, b be elements of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. If $a \Leftrightarrow b = true(Y)$, then $\forall_{a,P_1} G \Leftrightarrow \forall_{b,P_1} G = true(Y)$.

(28) Let a, b be elements of $Boolean^Y$, G be a subset of PARTITIONS(Y), and P_1 be a partition of Y. If $a \Leftrightarrow b = true(Y)$, then $\exists_{a,P_1} G \Leftrightarrow \exists_{b,P_1} G = true(Y)$.

REFERENCES

[1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[2] Shunichi Kobayashi and Kui Jia. A theory of Boolean valued functions and partitions. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/bvfunc_1.html.

[3] Shunichi Kobayashi and Yatsuka Nakamura. A theory of Boolean valued functions and quantifiers with respect to partitions. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/bvfunc_2.html.

[4] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/eqrel_1.html`.

[5] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[6] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/fraenkel.html`.

[7] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[8] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

[9] Edmund Woronowicz. Interpretation and satisfiability in the first order logic. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/valuat_1.html`.

[10] Edmund Woronowicz. Many-argument relations. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/margrel1.html`.

Received March 13, 1999

Published January 2, 2004