JOURNAL OF FORMALIZED MATHEMATICS
Volume 11, Released 1999, Published 2003
Inst. of Computer Science, Univ. of Białystok

Propositional Calculus for Boolean Valued Functions. Part I

Shunichi Kobayashi
Shinshu University
Nagano

Yatsuka Nakamura
Shinshu University
Nagano

Summary. In this paper, we have proved some elementary propositional calculus formulae for Boolean valued functions.

MML Identifier: BVFUNC_5.

WWW: http://mizar.org/JFM/Vol11/bvfunc_5.html

The articles [3], [5], [8], [7], [6], [1], [4], and [2] provide the notation and terminology for this paper.

In this paper Y denotes a non empty set.

We now state a number of propositions:

(1) For all elements a, b of $Boolean^Y$ holds $a = true(Y)$ and $b = true(Y)$ iff $a \wedge b = true(Y)$.

(2) For all elements a, b of $Boolean^Y$ such that $a = true(Y)$ and $a \Rightarrow b = true(Y)$ holds $b = true(Y)$.

(3) For all elements a, b of $Boolean^Y$ such that $a = true(Y)$ holds $a \vee b = true(Y)$.

(5)[1] For all elements a, b of $Boolean^Y$ such that $b = true(Y)$ holds $a \Rightarrow b = true(Y)$.

(6) For all elements a, b of $Boolean^Y$ such that $\neg a = true(Y)$ holds $a \Rightarrow b = true(Y)$.

(7) For every element a of $Boolean^Y$ holds $\neg(a \wedge \neg a) = true(Y)$.

(8) For every element a of $Boolean^Y$ holds $a \Rightarrow a = true(Y)$.

(9) For all elements a, b of $Boolean^Y$ holds $a \Rightarrow b = true(Y)$ iff $\neg b \Rightarrow \neg a = true(Y)$.

(10) For all elements a, b, c of $Boolean^Y$ such that $a \Rightarrow b = true(Y)$ and $b \Rightarrow c = true(Y)$ holds $a \Rightarrow c = true(Y)$.

(11) For all elements a, b of $Boolean^Y$ such that $a \Rightarrow b = true(Y)$ and $a \Rightarrow \neg b = true(Y)$ holds $\neg a = true(Y)$.

(12) For every element a of $Boolean^Y$ holds $\neg a \Rightarrow a \Rightarrow a = true(Y)$.

(13) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \Rightarrow \neg(b \wedge c) \Rightarrow \neg(a \wedge c) = true(Y)$.

(14) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \Rightarrow b \Rightarrow c \Rightarrow a \Rightarrow c = true(Y)$.

[1] The proposition (4) has been removed.

© Association of Mizar Users

(15) For all elements a, b, c of $Boolean^Y$ such that $a \Rightarrow b = true(Y)$ holds $b \Rightarrow c \Rightarrow a \Rightarrow c = true(Y)$.

(16) For all elements a, b of $Boolean^Y$ holds $b \Rightarrow a \Rightarrow b = true(Y)$.

(17) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \Rightarrow c \Rightarrow b \Rightarrow c = true(Y)$.

(18) For all elements a, b of $Boolean^Y$ holds $b \Rightarrow b \Rightarrow a \Rightarrow a = true(Y)$.

(19) For all elements a, b, c of $Boolean^Y$ holds $c \Rightarrow b \Rightarrow a \Rightarrow b \Rightarrow c \Rightarrow a = true(Y)$.

(20) For all elements a, b, c of $Boolean^Y$ holds $b \Rightarrow c \Rightarrow a \Rightarrow b \Rightarrow a \Rightarrow c = true(Y)$.

(21) For all elements a, b, c of $Boolean^Y$ holds $b \Rightarrow b \Rightarrow c \Rightarrow b \Rightarrow c = true(Y)$.

(22) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \Rightarrow c \Rightarrow a \Rightarrow b \Rightarrow a \Rightarrow c = true(Y)$.

(23) For all elements a, b of $Boolean^Y$ such that $a = true(Y)$ holds $a \Rightarrow b \Rightarrow b = true(Y)$.

(24) For all elements a, b, c of $Boolean^Y$ such that $c \Rightarrow b \Rightarrow a = true(Y)$ holds $b \Rightarrow c \Rightarrow a = true(Y)$.

(25) For all elements a, b, c of $Boolean^Y$ such that $c \Rightarrow b \Rightarrow a = true(Y)$ and $b = true(Y)$ holds $c \Rightarrow a = true(Y)$.

(26) For all elements a, b, c of $Boolean^Y$ such that $c \Rightarrow b \Rightarrow a = true(Y)$ and $b = true(Y)$ and $c = true(Y)$ holds $a = true(Y)$.

(27) For all elements b, c of $Boolean^Y$ such that $b \Rightarrow b \Rightarrow c = true(Y)$ holds $b \Rightarrow c = true(Y)$.

(28) For all elements a, b, c of $Boolean^Y$ such that $a \Rightarrow b \Rightarrow c = true(Y)$ holds $a \Rightarrow b \Rightarrow a \Rightarrow c = true(Y)$.

(29) For all elements a, b, c of $Boolean^Y$ such that $a \Rightarrow b \Rightarrow c = true(Y)$ and $a \Rightarrow b = true(Y)$ holds $a \Rightarrow c = true(Y)$.

(30) For all elements a, b, c of $Boolean^Y$ such that $a \Rightarrow b \Rightarrow c = true(Y)$ and $a \Rightarrow b = true(Y)$ and $a = true(Y)$ holds $c = true(Y)$.

(31) For all elements a, b, c, d of $Boolean^Y$ such that $a \Rightarrow b \Rightarrow c = true(Y)$ and $a \Rightarrow c \Rightarrow d = true(Y)$ holds $a \Rightarrow b \Rightarrow d = true(Y)$.

(32) For all elements a, b of $Boolean^Y$ holds $\neg a \Rightarrow \neg b \Rightarrow b \Rightarrow a = true(Y)$.

(33) For all elements a, b of $Boolean^Y$ holds $a \Rightarrow b \Rightarrow \neg b \Rightarrow \neg a = true(Y)$.

(34) For all elements a, b of $Boolean^Y$ holds $a \Rightarrow \neg b \Rightarrow b \Rightarrow \neg a = true(Y)$.

(35) For all elements a, b of $Boolean^Y$ holds $\neg a \Rightarrow b \Rightarrow \neg b \Rightarrow a = true(Y)$.

(36) For every element a of $Boolean^Y$ holds $a \Rightarrow \neg a \Rightarrow \neg a = true(Y)$.

(37) For all elements a, b of $Boolean^Y$ holds $\neg a \Rightarrow a \Rightarrow b = true(Y)$.

(38) For all elements a, b, c of $Boolean^Y$ holds $\neg(a \wedge b \wedge c) = \neg a \vee \neg b \vee \neg c$.

(39) For all elements a, b, c of $Boolean^Y$ holds $\neg(a \vee b \vee c) = \neg a \wedge \neg b \wedge \neg c$.

(40) For all elements a, b, c, d of $Boolean^Y$ holds $a \vee b \wedge c \wedge d = (a \vee b) \wedge (a \vee c) \wedge (a \vee d)$.

(41) For all elements a, b, c, d of $Boolean^Y$ holds $a \wedge (b \vee c \vee d) = a \wedge b \vee a \wedge c \vee a \wedge d$.

REFERENCES

[1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[2] Shunichi Kobayashi and Kui Jia. A theory of Boolean valued functions and partitions. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/bvfunc_1.html.

[3] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[4] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fraenkel.html.

[5] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[6] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[7] Edmund Woronowicz. Interpretation and satisfiability in the first order logic. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/valuat_1.html.

[8] Edmund Woronowicz. Many-argument relations. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/margrel1.html.

Received March 13, 1999

Published January 2, 2004