JOURNAL OF FORMALIZED MATHEMATICS
Volume 9, Released 1997, Published 2003
Inst. of Computer Science, Univ. of Białystok

Algebraic Operation on Subsets of Many Sorted Sets

Agnieszka Julia Marasik
Warsaw University
Białystok

MML Identifier: CLOSURE3.

WWW: http://mizar.org/JFM/Vol9/closure3.html

The articles [13], [5], [16], [12], [17], [2], [4], [3], [7], [6], [14], [15], [1], [10], [8], [9], and [11] provide the notation and terminology for this paper.

1. Preliminaries

Let S be a non empty 1-sorted structure. One can verify that the 1-sorted structure of S is non empty.

The following three propositions are true:

(1) For every non empty set I and for all many sorted sets M, N indexed by I holds $M + \cdot N = N$.

(2) Let I be a set, M, N be many sorted sets indexed by I, and F be a family of many sorted subsets indexed by M. If $N \in F$, then $\bigcap |{:}F{:}| \subseteq N$.

(3) Let S be a non void non empty many sorted signature, M_1 be a strict non-empty algebra over S, and F be a family of many sorted subsets indexed by the sorts of M_1. Suppose $F \subseteq \mathrm{SubSorts}(M_1)$. Let B be a subset of M_1. If $B = \bigcap |{:}F{:}|$, then B is operations closed.

2. Relationships between Subsets Families

Let I be a set, let M be a many sorted set indexed by I, let B be a family of many sorted subsets indexed by M, and let A be a family of many sorted subsets indexed by M. We say that A is finer than B if and only if:

(Def. 1) For every set a such that $a \in A$ there exists a set b such that $b \in B$ and $a \subseteq b$.

Let us note that the predicate A is finer than B is reflexive. We say that B is coarser than A if and only if:

(Def. 2) For every set b such that $b \in B$ there exists a set a such that $a \in A$ and $a \subseteq b$.

Let us note that the predicate B is coarser than A is reflexive.

One can prove the following two propositions:

(4) Let I be a set, M be a many sorted set indexed by I, and A, B, C be families of many sorted subsets indexed by M. If A is finer than B and B is finer than C, then A is finer than C.

(5) Let I be a set, M be a many sorted set indexed by I, and A, B, C be families of many sorted subsets indexed by M. If A is coarser than B and B is coarser than C, then A is coarser than C.

Let I be a non empty set and let M be a many sorted set indexed by I. The functor $\mathrm{supp}(M)$ yields a set and is defined as follows:

(Def. 3) $\mathrm{supp}(M) = \{x; x$ ranges over elements of I: $M(x) \neq \emptyset\}$.

The following propositions are true:

(6) For every non empty set I and for every non-empty many sorted set M indexed by I holds $M = \mathbf{0}_I {+\cdot} M \restriction \mathrm{supp}(M)$.

(7) Let I be a non empty set and M_2, M_3 be non-empty many sorted sets indexed by I. If $\mathrm{supp}(M_2) = \mathrm{supp}(M_3)$ and $M_2 \restriction \mathrm{supp}(M_2) = M_3 \restriction \mathrm{supp}(M_3)$, then $M_2 = M_3$.

(8) Let I be a non empty set, M be a many sorted set indexed by I, and x be an element of I. If $x \notin \mathrm{supp}(M)$, then $M(x) = \emptyset$.

(9) Let I be a non empty set, M be a many sorted set indexed by I, x be an element of $\mathrm{Bool}(M)$, i be an element of I, and y be a set. Suppose $y \in x(i)$. Then there exists an element a of $\mathrm{Bool}(M)$ such that $y \in a(i)$ and a is locally-finite and $\mathrm{supp}(a)$ is finite and $a \subseteq x$.

Let I be a set, let M be a many sorted set indexed by I, and let A be a family of many sorted subsets indexed by M. The functor $\mathrm{MSUnion}(A)$ yielding a many sorted subset indexed by M is defined as follows:

(Def. 4) For every set i such that $i \in I$ holds $(\mathrm{MSUnion}(A))(i) = \bigcup\{f(i); f$ ranges over elements of $\mathrm{Bool}(M)$: $f \in A\}$.

Let I be a set, let M be a many sorted set indexed by I, and let B be a non empty family of many sorted subsets indexed by M. We see that the element of B is a many sorted set indexed by I.

Let I be a set, let M be a many sorted set indexed by I, and let A be an empty family of many sorted subsets indexed by M. Observe that $\mathrm{MSUnion}(A)$ is empty yielding.

The following proposition is true

(10) Let I be a set, M be a many sorted set indexed by I, and A be a family of many sorted subsets indexed by M. Then $\mathrm{MSUnion}(A) = \bigcup|{:}A{:}|$.

Let I be a set, let M be a many sorted set indexed by I, and let A, B be families of many sorted subsets indexed by M. Then $A \cup B$ is a family of many sorted subsets indexed by M.

The following two propositions are true:

(11) Let I be a set, M be a many sorted set indexed by I, and A, B be families of many sorted subsets indexed by M. Then $\mathrm{MSUnion}(A \cup B) = \mathrm{MSUnion}(A) \cup \mathrm{MSUnion}(B)$.

(12) Let I be a set, M be a many sorted set indexed by I, and A, B be families of many sorted subsets indexed by M. If $A \subseteq B$, then $\mathrm{MSUnion}(A) \subseteq \mathrm{MSUnion}(B)$.

Let I be a set, let M be a many sorted set indexed by I, and let A, B be families of many sorted subsets indexed by M. Then $A \cap B$ is a family of many sorted subsets indexed by M.

One can prove the following two propositions:

(13) Let I be a set, M be a many sorted set indexed by I, and A, B be families of many sorted subsets indexed by M. Then $\mathrm{MSUnion}(A \cap B) \subseteq \mathrm{MSUnion}(A) \cap \mathrm{MSUnion}(B)$.

(14) Let I be a set, M be a many sorted set indexed by I, and A_1 be a set. Suppose that for every set x such that $x \in A_1$ holds x is a family of many sorted subsets indexed by M. Let A, B be families of many sorted subsets indexed by M. Suppose $B = \{\mathrm{MSUnion}(X); X$ ranges over families of many sorted subsets indexed by M: $X \in A_1\}$ and $A = \bigcup A_1$. Then $\mathrm{MSUnion}(B) = \mathrm{MSUnion}(A)$.

3. Algebraic Operation on Subsets of Many Sorted Sets

Let I be a non empty set, let M be a many sorted set indexed by I, and let S be a set operation in M. We say that S is algebraic if and only if the condition (Def. 5) is satisfied.

(Def. 5) Let x be an element of $\mathrm{Bool}(M)$. Suppose $x = S(x)$. Then there exists a family A of many sorted subsets indexed by M such that $A = \{S(a); a$ ranges over elements of $\mathrm{Bool}(M)$: a is locally-finite $\wedge$ $\mathrm{supp}(a)$is finite $\wedge$ $a \subseteq x\}$ and $x = \mathrm{MSUnion}(A)$.

Let I be a non empty set and let M be a many sorted set indexed by I. Note that there exists a set operation in M which is algebraic, reflexive, monotonic, and idempotent.

Let S be a non empty 1-sorted structure and let I_1 be a closure system of S. We say that I_1 is algebraic if and only if:

(Def. 6) $\mathrm{ClOp}(I_1)$ is algebraic.

Let S be a non void non empty many sorted signature and let M_1 be a non-empty algebra over S. The functor $\mathrm{SubAlgCl}(M_1)$ yielding a strict closure system structure over the 1-sorted structure of S is defined by:

(Def. 7) The sorts of $\mathrm{SubAlgCl}(M_1)$ = the sorts of M_1 and the family of $\mathrm{SubAlgCl}(M_1) = \mathrm{SubSorts}(M_1)$.

We now state the proposition

(16)[1] Let S be a non void non empty many sorted signature and M_1 be a strict non-empty algebra over S. Then $\mathrm{SubSorts}(M_1)$ is an absolutely-multiplicative family of many sorted subsets indexed by the sorts of M_1.

Let S be a non void non empty many sorted signature and let M_1 be a strict non-empty algebra over S. One can check that $\mathrm{SubAlgCl}(M_1)$ is absolutely-multiplicative.

Let S be a non void non empty many sorted signature and let M_1 be a strict non-empty algebra over S. Note that $\mathrm{SubAlgCl}(M_1)$ is algebraic.

References

[1] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_2.html.

[2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[4] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/partfun1.html.

[5] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[6] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.

[7] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[8] Artur Korniłowicz. Certain facts about families of subsets of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/mssubfam.html.

[9] Artur Korniłowicz. Definitions and basic properties of boolean and union of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/mboolean.html.

[10] Artur Korniłowicz. On the closure operator and the closure system of many sorted sets. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/closure2.html.

[11] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pre_circ.html.

[12] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.

[1] The proposition (15) has been removed.

[13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[14] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/pboole.html`.

[15] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. `http://mizar.org/JFM/Vol6/msualg_1.html`.

[16] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[17] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

Received June 23, 1997

Published January 2, 2004