JOURNAL OF FORMALIZED MATHEMATICS
Volume 8, Released 1996, Published 2003
Inst. of Computer Science, Univ. of Białystok

Components and Unions of Components

Yatsuka Nakamura
Shinshu University
Nagano

Andrzej Trybulec
Warsaw University
Białystok

Summary. First, we generalized **skl** function for a subset of topological spaces the value of which is the component including the set. Second, we introduced a concept of union of components a family of which has good algebraic properties. At the end, we discuss relationship between connectivity of a set as a subset in the whole space and as a subset of a subspace.

MML Identifier: `CONNSP_3`.

WWW: `http://mizar.org/JFM/Vol8/connsp_3.html`

The articles [5], [1], [3], [4], and [2] provide the notation and terminology for this paper.

1. The Component of a Subset in a Topological Space

In this paper G_1 is a non empty topological space.

Let G_1 be a topological structure and let V be a subset of G_1. The functor Component(V) yielding a subset of G_1 is defined by the condition (Def. 1).

(Def. 1) There exists a family F of subsets of G_1 such that for every subset A of G_1 holds $A \in F$ iff A is connected and $V \subseteq A$ and $\bigcup F = \mathrm{Component}(V)$.

One can prove the following propositions:

(1) Let G_1 be a topological space and V be a subset of G_1. If there exists a subset A of G_1 such that A is connected and $V \subseteq A$, then $V \subseteq \mathrm{Component}(V)$.

(2) Let G_1 be a topological space and V be a subset of G_1. If it is not true that there exists a subset A of G_1 such that A is connected and $V \subseteq A$, then $\mathrm{Component}(V) = \emptyset$.

(3) $\mathrm{Component}(\emptyset_{(G_1)})$ = the carrier of G_1.

(4) For every subset V of G_1 such that V is connected holds $\mathrm{Component}(V) \neq \emptyset$.

(5) For every topological space G_1 and for every subset V of G_1 such that V is connected and $V \neq \emptyset$ holds Component(V) is connected.

(6) For all subsets V, C of G_1 such that V is connected and C is connected holds if $\mathrm{Component}(V) \subseteq C$, then $C = \mathrm{Component}(V)$.

(7) For every subset A of G_1 such that A is a component of G_1 holds $\mathrm{Component}(A) = A$.

(8) Let A be a subset of G_1. Then A is a component of G_1 if and only if there exists a subset V of G_1 such that V is connected and $V \neq \emptyset$ and $A = \mathrm{Component}(V)$.

(9) For every subset V of G_1 such that V is connected and $V \neq \emptyset$ holds Component(V) is a component of G_1.

(10) For all subsets A, V of G_1 such that A is a component of G_1 and V is connected and $V \subseteq A$ and $V \neq \emptyset$ holds $A =$ Component(V).

(11) For every subset V of G_1 such that V is connected and $V \neq \emptyset$ holds Component(Component$(V)) =$ Component(V).

(12) For all subsets A, B of G_1 such that A is connected and B is connected and $A \neq \emptyset$ and $A \subseteq B$ holds Component$(A) =$ Component(B).

(13) For all subsets A, B of G_1 such that A is connected and B is connected and $A \neq \emptyset$ and $A \subseteq B$ holds $B \subseteq$ Component(A).

(14) Let A be a subset of G_1 and B be a subset of G_1. If A is connected and $A \cup B$ is connected and $A \neq \emptyset$, then $A \cup B \subseteq$ Component(A).

(15) For every subset A of G_1 and for every point p of G_1 such that A is connected and $p \in A$ holds Component$(p) =$ Component(A).

(16) Let A, B be subsets of G_1. Suppose A is connected and B is connected and A meets B. Then $A \cup B \subseteq$ Component(A) and $A \cup B \subseteq$ Component(B) and $A \subseteq$ Component(B) and $B \subseteq$ Component(A).

(17) For every subset A of G_1 such that A is connected and $A \neq \emptyset$ holds $\overline{A} \subseteq$ Component(A).

(18) Let A, B be subsets of G_1. Suppose A is a component of G_1 and B is connected and $B \neq \emptyset$ and A misses B. Then A misses Component(B).

2. On Unions of Components

Let G_1 be a topological structure. A subset of G_1 is called a union of components of G_1 if it satisfies the condition (Def. 2).

(Def. 2) There exists a family F of subsets of G_1 such that for every subset B of G_1 such that $B \in F$ holds B is a component of G_1 and it $= \bigcup F$.

The following propositions are true:

(19) $\emptyset_{(G_1)}$ is a union of components of G_1.

(20) For every subset A of G_1 such that $A =$ the carrier of G_1 holds A is a union of components of G_1.

(21) Let A be a subset of G_1 and p be a point of G_1. If $p \in A$ and A is a union of components of G_1, then Component$(p) \subseteq A$.

(22) Let A, B be subsets of G_1. Suppose A is a union of components of G_1 and B is a union of components of G_1. Then $A \cup B$ is a union of components of G_1 and $A \cap B$ is a union of components of G_1.

(23) Let F_1 be a family of subsets of G_1. Suppose that for every subset A of G_1 such that $A \in F_1$ holds A is a union of components of G_1. Then $\bigcup F_1$ is a union of components of G_1.

(24) Let F_1 be a family of subsets of G_1. Suppose that for every subset A of G_1 such that $A \in F_1$ holds A is a union of components of G_1. Then $\bigcap F_1$ is a union of components of G_1.

(25) Let A, B be subsets of G_1. Suppose A is a union of components of G_1 and B is a union of components of G_1. Then $A \setminus B$ is a union of components of G_1.

3. Operations Down and Up

Let G_1 be a topological structure, let B be a subset of G_1, and let p be a point of G_1. Let us assume that $p \in B$. The functor $\mathrm{Down}(p,B)$ yields a point of $G_1 \restriction B$ and is defined as follows:

(Def. 3) $\mathrm{Down}(p,B) = p$.

Let G_1 be a topological structure, let B be a subset of G_1, and let p be a point of $G_1 \restriction B$. Let us assume that $B \neq \emptyset$. The functor $\mathrm{Up}(p)$ yielding a point of G_1 is defined as follows:

(Def. 4) $\mathrm{Up}(p) = p$.

Let G_1 be a topological structure and let V, B be subsets of G_1. The functor $\mathrm{Down}(V,B)$ yielding a subset of $G_1 \restriction B$ is defined by:

(Def. 5) $\mathrm{Down}(V,B) = V \cap B$.

Let G_1 be a topological structure, let B be a subset of G_1, and let V be a subset of $G_1 \restriction B$. The functor $\mathrm{Up}(V)$ yields a subset of G_1 and is defined by:

(Def. 6) $\mathrm{Up}(V) = V$.

Let G_1 be a topological structure, let B be a subset of G_1, and let p be a point of G_1. Let us assume that $p \in B$. The functor $\mathrm{skl}(p,B)$ yielding a subset of G_1 is defined as follows:

(Def. 7) For every point q of $G_1 \restriction B$ such that $q = p$ holds $\mathrm{skl}(p,B) = \mathrm{Component}(q)$.

The following propositions are true:

(26) For every subset B of G_1 and for every point p of G_1 such that $p \in B$ holds $\mathrm{skl}(p,B) \neq \emptyset$.

(27) For every subset B of G_1 and for every point p of G_1 such that $p \in B$ holds $\mathrm{skl}(p,B) = \mathrm{Component}(\mathrm{Down}(p,B))$.

(28) For all subsets V, B of G_1 such that $V \subseteq B$ holds $\mathrm{Down}(V,B) = V$.

(29) For every topological space G_1 and for all subsets V, B of G_1 such that V is open holds $\mathrm{Down}(V,B)$ is open.

(30) For all subsets V, B of G_1 such that $V \subseteq B$ holds $\overline{\mathrm{Down}(V,B)} = \overline{V} \cap B$.

(31) For every subset B of G_1 and for every subset V of $G_1 \restriction B$ holds $\overline{V} = \overline{\mathrm{Up}(V)} \cap B$.

(32) For all subsets V, B of G_1 such that $V \subseteq B$ holds $\overline{\mathrm{Down}(V,B)} \subseteq \overline{V}$.

(33) For every subset B of G_1 and for every subset V of $G_1 \restriction B$ such that $V \subseteq B$ holds $\mathrm{Down}(\mathrm{Up}(V),B) = V$.

(34) Let G_1 be a topological space, V, B be subsets of G_1, and W be a subset of $G_1 \restriction B$. If $V = W$ and W is connected, then V is connected.

(35) For every subset B of G_1 and for every point p of G_1 such that $p \in B$ holds $\mathrm{skl}(p,B)$ is connected.

References

[1] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/zfmisc_1.html`.

[2] Beata Padlewska. Connected spaces. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/connsp_1.html`.

[3] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/setfam_1.html`.

[4] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/pre_topc.html`.

[5] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

Received February 5, 1996

Published January 2, 2004