JOURNAL OF FORMALIZED MATHEMATICS
Volume 7, Released 1995, Published 2003
Inst. of Computer Science, Univ. of Białystok

On the Decomposition of Finite Sequences

Andrzej Trybulec
Warsaw University
Białystok

MML Identifier: FINSEQ_6.

WWW: http://mizar.org/JFM/Vol7/finseq_6.html

The articles [10], [13], [2], [3], [11], [1], [14], [15], [6], [4], [12], [9], [7], [5], and [8] provide the notation and terminology for this paper.

1. Preliminaries

In this paper x, y, z denote sets.

Let us consider x, y, z. One can check that $\langle x, y, z \rangle$ is non trivial.

Let f be a non empty finite sequence. One can check that $\mathrm{Rev}(f)$ is non empty.

2. Decomposing a finite sequence

We adopt the following rules: f, f_1, f_2, f_3 are finite sequences, p, p_1, p_2, p_3 are sets, and i, k are natural numbers.

The following propositions are true:

(3)[1] For every set X and for every i such that $X \subseteq \mathrm{Seg}\, i$ and $1 \in X$ holds $(\mathrm{Sgm} X)(1) = 1$.

(4) For every finite sequence f such that $k \in \mathrm{dom} f$ and for every i such that $1 \leq i$ and $i < k$ holds $f(i) \neq f(k)$ holds $f(k) \leftarrow\!\!\!\!\!\!\!\!\!\;\;\;\;\;\; f = k$.

(5) $\langle p_1, p_2 \rangle \restriction \mathrm{Seg}\, 1 = \langle p_1 \rangle$.

(6) $\langle p_1, p_2, p_3 \rangle \restriction \mathrm{Seg}\, 1 = \langle p_1 \rangle$.

(7) $\langle p_1, p_2, p_3 \rangle \restriction \mathrm{Seg}\, 2 = \langle p_1, p_2 \rangle$.

(8) If $p \in \mathrm{rng} f_1$, then $p \leftarrow (f_1 \frown f_2) = p \leftarrow f_1$.

(9) If $p \in \mathrm{rng} f_2 \setminus \mathrm{rng} f_1$, then $p \leftarrow (f_1 \frown f_2) = \mathrm{len} f_1 + p \leftarrow f_2$.

(10) If $p \in \mathrm{rng} f_1$, then $f_1 \frown f_2 \rightarrow p = (f_1 \rightarrow p) \frown f_2$.

(11) If $p \in \mathrm{rng} f_2 \setminus \mathrm{rng} f_1$, then $f_1 \frown f_2 \rightarrow p = f_2 \rightarrow p$.

(12) $f_1 \subseteq f_1 \frown f_2$.

(13) For every set A such that $A \subseteq \mathrm{dom} f_1$ holds $(f_1 \frown f_2) \restriction A = f_1 \restriction A$.

[1] The propositions (1) and (2) have been removed.

(14) If $p \in \operatorname{rng} f_1$, then $f_1 \frown f_2 \leftarrow p = f_1 \leftarrow p$.

Let us consider f_1 and let i be a natural number. Note that $f_1 \restriction \operatorname{Seg} i$ is finite sequence-like.
The following propositions are true:

(15) If $f_1 \subseteq f_2$, then $f_3 \frown f_1 \subseteq f_3 \frown f_2$.

(16) $(f_1 \frown f_2) \restriction \operatorname{Seg}(\operatorname{len} f_1 + i) = f_1 \frown (f_2 \restriction \operatorname{Seg} i)$.

(17) If $p \in \operatorname{rng} f_2 \setminus \operatorname{rng} f_1$, then $f_1 \frown f_2 \leftarrow p = f_1 \frown (f_2 \leftarrow p)$.

(19)[2] If $f_1 \frown f_2$ yields p just once, then $p \in \operatorname{rng} f_1 \dot{-} \operatorname{rng} f_2$.

(20) If $f_1 \frown f_2$ yields p just once and $p \in \operatorname{rng} f_1$, then f_1 yields p just once.

(21) If $\operatorname{rng} f$ is non trivial, then f is non trivial.

(22) $p \hookleftarrow \langle p \rangle = 1$.

(23) $p_1 \hookleftarrow \langle p_1, p_2 \rangle = 1$.

(24) If $p_1 \neq p_2$, then $p_2 \hookleftarrow \langle p_1, p_2 \rangle = 2$.

(25) $p_1 \hookleftarrow \langle p_1, p_2, p_3 \rangle = 1$.

(26) If $p_1 \neq p_2$, then $p_2 \hookleftarrow \langle p_1, p_2, p_3 \rangle = 2$.

(27) If $p_1 \neq p_3$ and $p_2 \neq p_3$, then $p_3 \hookleftarrow \langle p_1, p_2, p_3 \rangle = 3$.

(28) For every finite sequence f holds $\operatorname{Rev}(\langle p \rangle \frown f) = (\operatorname{Rev}(f)) \frown \langle p \rangle$.

(29) For every finite sequence f holds $\operatorname{Rev}(\operatorname{Rev}(f)) = f$.

(30) If $x \neq y$, then $\langle x, y \rangle \leftarrow y = \langle x \rangle$.

(31) If $x \neq y$, then $\langle x, y, z \rangle \leftarrow y = \langle x \rangle$.

(32) If $x \neq z$ and $y \neq z$, then $\langle x, y, z \rangle \leftarrow z = \langle x, y \rangle$.

(33) $\langle x, y \rangle \rightarrow x = \langle y \rangle$.

(34) If $x \neq y$, then $\langle x, y, z \rangle \rightarrow y = \langle z \rangle$.

(35) $\langle x, y, z \rangle \rightarrow x = \langle y, z \rangle$.

(36) $\langle z \rangle \rightarrow z = \emptyset$ and $\langle z \rangle \leftarrow z = \emptyset$.

(37) If $x \neq y$, then $\langle x, y \rangle \rightarrow y = \emptyset$.

(38) If $x \neq z$ and $y \neq z$, then $\langle x, y, z \rangle \rightarrow z = \emptyset$.

(39) If $x \in \operatorname{rng} f$ and $y \in \operatorname{rng}(f \leftarrow x)$, then $(f \leftarrow x) \leftarrow y = f \leftarrow y$.

(40) If $x \notin \operatorname{rng} f_1$, then $x \hookleftarrow (f_1 \frown \langle x \rangle \frown f_2) = \operatorname{len} f_1 + 1$.

(41) If f yields x just once, then $x \hookleftarrow f + x \hookleftarrow \operatorname{Rev}(f) = \operatorname{len} f + 1$.

(42) If f yields x just once, then $\operatorname{Rev}(f \leftarrow x) = \operatorname{Rev}(f) \rightarrow x$.

(43) If f yields x just once, then $\operatorname{Rev}(f)$ yields x just once.

[2] The proposition (18) has been removed.

3. Finite sequences with elements from a non empty set

We adopt the following convention: D denotes a non empty set, p, p_1, p_2, p_3 denote elements of D, and f, f_1, f_2 denote finite sequences of elements of D.

One can prove the following propositions:

(44) If $p \in \operatorname{rng} f$, then $f -: p = (f \leftarrow p) ^\frown \langle p \rangle$.

(45) If $p \in \operatorname{rng} f$, then $f :- p = \langle p \rangle ^\frown (f \rightarrow p)$.

(46) If $f \neq \emptyset$, then $f_1 \in \operatorname{rng} f$.

(47) If $f \neq \emptyset$, then $f_1 \looparrowleft f = 1$.

(48) If $f \neq \emptyset$ and $f_1 = p$, then $f -: p = \langle p \rangle$ and $f :- p = f$.

(49) $(\langle p_1 \rangle ^\frown f)_{\downharpoonleft 1} = f$.

(50) $\langle p_1, p_2 \rangle_{\downharpoonleft 1} = \langle p_2 \rangle$.

(51) $\langle p_1, p_2, p_3 \rangle_{\downharpoonleft 1} = \langle p_2, p_3 \rangle$.

(52) If $k \in \operatorname{dom} f$ and for every i such that $1 \leq i$ and $i < k$ holds $f_i \neq f_k$, then $f_k \looparrowleft f = k$.

(53) If $p_1 \neq p_2$, then $\langle p_1, p_2 \rangle -: p_2 = \langle p_1, p_2 \rangle$.

(54) If $p_1 \neq p_2$, then $\langle p_1, p_2, p_3 \rangle -: p_2 = \langle p_1, p_2 \rangle$.

(55) If $p_1 \neq p_3$ and $p_2 \neq p_3$, then $\langle p_1, p_2, p_3 \rangle -: p_3 = \langle p_1, p_2, p_3 \rangle$.

(56) $\langle p \rangle :- p = \langle p \rangle$ and $\langle p \rangle -: p = \langle p \rangle$.

(57) If $p_1 \neq p_2$, then $\langle p_1, p_2 \rangle :- p_2 = \langle p_2 \rangle$.

(58) If $p_1 \neq p_2$, then $\langle p_1, p_2, p_3 \rangle :- p_2 = \langle p_2, p_3 \rangle$.

(59) If $p_1 \neq p_3$ and $p_2 \neq p_3$, then $\langle p_1, p_2, p_3 \rangle :- p_3 = \langle p_3 \rangle$.

(61)[3] If $p \in \operatorname{rng} f$ and $p \looparrowleft f > k$, then $p \looparrowleft f = k + p \looparrowleft (f_{\downharpoonleft k})$.

(62) If $p \in \operatorname{rng} f$ and $p \looparrowleft f > k$, then $p \in \operatorname{rng}(f_{\downharpoonleft k})$.

(63) If $k < i$ and $i \in \operatorname{dom} f$, then $f_i \in \operatorname{rng}(f_{\downharpoonleft k})$.

(64) If $p \in \operatorname{rng} f$ and $p \looparrowleft f > k$, then $f_{\downharpoonleft k} -: p = (f -: p)_{\downharpoonleft k}$.

(65) If $p \in \operatorname{rng} f$ and $p \looparrowleft f \neq 1$, then $f_{\downharpoonleft 1} -: p = (f -: p)_{\downharpoonleft 1}$.

(66) $p \in \operatorname{rng}(f :- p)$.

(67) If $x \in \operatorname{rng} f$ and $p \in \operatorname{rng} f$ and $x \looparrowleft f \geq p \looparrowleft f$, then $x \in \operatorname{rng}(f :- p)$.

(68) If $p \in \operatorname{rng} f$ and $k \leq \operatorname{len} f$ and $k \geq p \looparrowleft f$, then $f_k \in \operatorname{rng}(f :- p)$.

(69) If $p \in \operatorname{rng} f_1$, then $f_1 ^\frown f_2 :- p = (f_1 :- p) ^\frown f_2$.

(70) If $p \in \operatorname{rng} f_2 \setminus \operatorname{rng} f_1$, then $f_1 ^\frown f_2 :- p = f_2 :- p$.

(71) If $p \in \operatorname{rng} f_1$, then $f_1 ^\frown f_2 -: p = f_1 -: p$.

(72) If $p \in \operatorname{rng} f_2 \setminus \operatorname{rng} f_1$, then $f_1 ^\frown f_2 -: p = f_1 ^\frown (f_2 -: p)$.

(73) $f :- p :- p = f :- p$.

(74) If $p_1 \in \operatorname{rng} f$ and $p_2 \in \operatorname{rng} f \setminus \operatorname{rng}(f -: p_1)$, then $f \rightarrow p_2 = (f \rightarrow p_1) \rightarrow p_2$.

[3] The proposition (60) has been removed.

(75) If $p \in \operatorname{rng} f$, then $\operatorname{rng} f = \operatorname{rng}(f -: p) \cup \operatorname{rng}(f :- p)$.

(76) If $p_1 \in \operatorname{rng} f$ and $p_2 \in \operatorname{rng} f \setminus \operatorname{rng}(f -: p_1)$, then $f :- p_1 :- p_2 = f :- p_2$.

(77) If $p \in \operatorname{rng} f$, then $p \leftrightarrow (f -: p) = p \leftrightarrow f$.

(78) $f \restriction i \restriction i = f \restriction i$.

(79) If $p \in \operatorname{rng} f$, then $f -: p -: p = f -: p$.

(80) If $p_1 \in \operatorname{rng} f$ and $p_2 \in \operatorname{rng}(f -: p_1)$, then $f -: p_1 -: p_2 = f -: p_2$.

(81) If $p \in \operatorname{rng} f$, then $(f -: p) \frown ((f :- p)_{\downharpoonright 1}) = f$.

(82) If $f_1 \neq \emptyset$, then $(f_1 \frown f_2)_{\downharpoonright 1} = ((f_1)_{\downharpoonright 1}) \frown f_2$.

(83) If $p_2 \in \operatorname{rng} f$ and $p_2 \leftrightarrow f \neq 1$, then $p_2 \in \operatorname{rng}(f_{\downharpoonright 1})$.

(84) $p \leftrightarrow (f :- p) = 1$.

(86)[4] $(\varepsilon_D)_{\downharpoonright k} = \varepsilon_D$.

(87) $f_{\downharpoonright i+k} = (f_{\downharpoonright i})_{\downharpoonright k}$.

(88) If $p \in \operatorname{rng} f$ and $p \leftrightarrow f > k$, then $f_{\downharpoonright k} :- p = f :- p$.

(89) If $p \in \operatorname{rng} f$ and $p \leftrightarrow f \neq 1$, then $f_{\downharpoonright 1} :- p = f :- p$.

(90) If $i + k = \operatorname{len} f$, then $\operatorname{Rev}(f_{\downharpoonright k}) = \operatorname{Rev}(f) \restriction i$.

(91) If $i + k = \operatorname{len} f$, then $\operatorname{Rev}(f \restriction k) = (\operatorname{Rev}(f))_{\downharpoonright i}$.

(92) If f yields p just once, then $\operatorname{Rev}(f \rightarrow p) = \operatorname{Rev}(f) \leftarrow p$.

(93) If f yields p just once, then $\operatorname{Rev}(f :- p) = \operatorname{Rev}(f) -: p$.

(94) If f yields p just once, then $\operatorname{Rev}(f -: p) = \operatorname{Rev}(f) :- p$.

4. Rotating a finite sequence

Let D be a non empty set and let I_1 be a finite sequence of elements of D. We say that I_1 is circular if and only if:

(Def. 1) $(I_1)_1 = (I_1)_{\operatorname{len} I_1}$.

Let us consider D, f, p. The functor $f \circlearrowleft p$ yielding a finite sequence of elements of D is defined as follows:

(Def. 2) $$f \circlearrowleft p = \begin{cases} (f :- p) \frown ((f -: p)_{\downharpoonright 1}), & \text{if } p \in \operatorname{rng} f, \\ f, & \text{otherwise.} \end{cases}$$

Let us consider D, let f be a non empty finite sequence of elements of D, and let p be an element of D. Observe that $f \circlearrowleft p$ is non empty.

Let us consider D. Observe that there exists a finite sequence of elements of D which is circular, non empty, and trivial and there exists a finite sequence of elements of D which is circular, non empty, and non trivial.

One can prove the following proposition

(95) $f \circlearrowleft f_1 = f$.

Let us consider D, p and let f be a circular non empty finite sequence of elements of D. Note that $f \circlearrowleft p$ is circular.

We now state a number of propositions:

[4] The proposition (85) has been removed.

(96) If f is circular and $p \in \operatorname{rng} f$, then $\operatorname{rng}(f \circlearrowleft p) = \operatorname{rng} f$.

(97) If $p \in \operatorname{rng} f$, then $p \in \operatorname{rng}(f \circlearrowleft p)$.

(98) If $p \in \operatorname{rng} f$, then $(f \circlearrowleft p)_1 = p$.

(99) $(f \circlearrowleft p) \circlearrowleft p = f \circlearrowleft p$.

(100) $\langle p \rangle \circlearrowleft p = \langle p \rangle$.

(101) $\langle p_1, p_2 \rangle \circlearrowleft p_1 = \langle p_1, p_2 \rangle$.

(102) $\langle p_1, p_2 \rangle \circlearrowleft p_2 = \langle p_2, p_2 \rangle$.

(103) $\langle p_1, p_2, p_3 \rangle \circlearrowleft p_1 = \langle p_1, p_2, p_3 \rangle$.

(104) If $p_1 \neq p_2$, then $\langle p_1, p_2, p_3 \rangle \circlearrowleft p_2 = \langle p_2, p_3, p_2 \rangle$.

(105) If $p_2 \neq p_3$, then $\langle p_1, p_2, p_3 \rangle \circlearrowleft p_3 = \langle p_3, p_2, p_3 \rangle$.

(106) For every circular non trivial finite sequence f of elements of D holds $\operatorname{rng}(f_{\lfloor 1}) = \operatorname{rng} f$.

(107) $\operatorname{rng}(f_{\lfloor 1}) \subseteq \operatorname{rng}(f \circlearrowleft p)$.

(108) If $p_2 \in \operatorname{rng} f \setminus \operatorname{rng}(f -: p_1)$, then $(f \circlearrowleft p_1) \circlearrowleft p_2 = f \circlearrowleft p_2$.

(109) If $p_2 \leftarrow f \neq 1$ and $p_2 \in \operatorname{rng} f \setminus \operatorname{rng}(f :- p_1)$, then $(f \circlearrowleft p_1) \circlearrowleft p_2 = f \circlearrowleft p_2$.

(110) If $p_2 \in \operatorname{rng}(f_{\lfloor 1})$ and f yields p_2 just once, then $(f \circlearrowleft p_1) \circlearrowleft p_2 = f \circlearrowleft p_2$.

(111) If f is circular and f yields p_2 just once, then $(f \circlearrowleft p_1) \circlearrowleft p_2 = f \circlearrowleft p_2$.

(112) If f is circular and f yields p just once, then $\operatorname{Rev}(f \circlearrowleft p) = \operatorname{Rev}(f) \circlearrowleft p$.

REFERENCES

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/nat_1.html`.

[2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/ordinal1.html`.

[3] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/ordinal2.html`.

[4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/finseq_1.html`.

[5] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/realset1.html`.

[6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_1.html`.

[7] Czesław Byliński. Some properties of restrictions of finite sequences. *Journal of Formalized Mathematics*, 7, 1995. `http://mizar.org/JFM/Vol7/finseq_5.html`.

[8] Agata Darmochwał and Yatsuka Nakamura. The topological space $\mathcal{E}^2_T$. Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. `http://mizar.org/JFM/Vol3/topreal1.html`.

[9] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/rfinseq.html`.

[10] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[11] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. `http://mizar.org/JFM/Addenda/numbers.html`.

[12] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/finseq_4.html`.

[13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[14] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

[15] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relset_1.html`.

Received May 24, 1995

Published January 2, 2004