JOURNAL OF FORMALIZED MATHEMATICS
Volume 1, Released 1989, Published 2003
Inst. of Computer Science, Univ. of Białystok

Finite Sets

Agata Darmochwał
Warsaw University
Białystok

Summary. The article contains the definition of a finite set based on the notion of finite sequence. Some theorems about properties of finite sets and finite families of sets are proved.

MML Identifier: FINSET_1.

WWW: http://mizar.org/JFM/Vol1/finset_1.html

The articles [6], [4], [7], [8], [3], [5], [1], and [2] provide the notation and terminology for this paper.

Let I_1 be a set. We say that I_1 is finite if and only if:

(Def. 1) There exists a function p such that $\operatorname{rng} p = I_1$ and $\operatorname{dom} p \in \omega$.

We introduce I_1 is infinite as an antonym of I_1 is finite.

We adopt the following convention: A, B, C, D, X, Y are sets and f is a function.

Let us mention that there exists a set which is non empty and finite.

One can verify that every set which is empty is also finite.

Let X be a set. Observe that there exists a subset of X which is empty and finite.

The scheme *OLambdaC* deals with a set $\mathcal{A}$, a unary functor $\mathcal{F}$ yielding a set, a unary functor $\mathcal{G}$ yielding a set, and a unary predicate $\mathcal{P}$, and states that:

> There exists a function f such that $\operatorname{dom} f = \mathcal{A}$ and for every ordinal number x such that $x \in \mathcal{A}$ holds if $\mathcal{P}[x]$, then $f(x) = \mathcal{F}(x)$ and if not $\mathcal{P}[x]$, then $f(x) = \mathcal{G}(x)$

for all values of the parameters.

Let x be a set. One can verify that $\{x\}$ is finite.

Let X be a non empty set. Observe that there exists a subset of X which is non empty and finite.

Let x, y be sets. One can verify that $\{x,y\}$ is finite.

Let x, y, z be sets. One can check that $\{x,y,z\}$ is finite.

Let x_1, x_2, x_3, x_4 be sets. One can verify that $\{x_1,x_2,x_3,x_4\}$ is finite.

Let x_1, x_2, x_3, x_4, x_5 be sets. Observe that $\{x_1,x_2,x_3,x_4,x_5\}$ is finite.

Let x_1, x_2, x_3, x_4, x_5, x_6 be sets. Observe that $\{x_1,x_2,x_3,x_4,x_5,x_6\}$ is finite.

Let x_1, x_2, x_3, x_4, x_5, x_6, x_7 be sets. Observe that $\{x_1,x_2,x_3,x_4,x_5,x_6,x_7\}$ is finite.

Let x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8 be sets. One can verify that $\{x_1,x_2,x_3,x_4,x_5,x_6,x_7,x_8\}$ is finite.

Let B be a finite set. Note that every subset of B is finite.

Let X, Y be finite sets. Note that $X \cup Y$ is finite.

One can prove the following two propositions:

(13)[1] If $A \subseteq B$ and B is finite, then A is finite.

(14) If A is finite and B is finite, then $A \cup B$ is finite.

[1] The propositions (1)–(12) have been removed.

© Association of Mizar Users

Let A be a set and let B be a finite set. One can check that $A \cap B$ is finite.
Let A be a finite set and let B be a set. Observe that $A \cap B$ is finite and $A \setminus B$ is finite.
Next we state three propositions:

(15) If A is finite, then $A \cap B$ is finite.

(16) If A is finite, then $A \setminus B$ is finite.

(17) If A is finite, then $f^{\circ}A$ is finite.

Let f be a function and let A be a finite set. Observe that $f^{\circ}A$ is finite.
One can prove the following proposition

(18) Suppose A is finite. Let X be a family of subsets of A. Suppose $X \neq \emptyset$. Then there exists a set x such that $x \in X$ and for every set B such that $B \in X$ holds if $x \subseteq B$, then $B = x$.

The scheme *Finite* deals with a set $\mathcal{A}$ and a unary predicate $\mathcal{P}$, and states that:
$\mathcal{P}[\mathcal{A}]$
provided the following requirements are met:

- $\mathcal{A}$ is finite,
- $\mathcal{P}[\emptyset]$, and
- For all sets x, B such that $x \in \mathcal{A}$ and $B \subseteq \mathcal{A}$ and $\mathcal{P}[B]$ holds $\mathcal{P}[B \cup \{x\}]$.

One can prove the following proposition

(19) If A is finite and B is finite, then $[:A, B:]$ is finite.

Let A, B be finite sets. Note that $[:A, B:]$ is finite.
We now state the proposition

(20) If A is finite and B is finite and C is finite, then $[:A, B, C:]$ is finite.

Let A, B, C be finite sets. One can check that $[:A, B, C:]$ is finite.
Next we state the proposition

(21) If A is finite and B is finite and C is finite and D is finite, then $[:A, B, C, D:]$ is finite.

Let A, B, C, D be finite sets. Observe that $[:A, B, C, D:]$ is finite.
The following propositions are true:

(22) If $B \neq \emptyset$ and $[:A, B:]$ is finite, then A is finite.

(23) If $A \neq \emptyset$ and $[:A, B:]$ is finite, then B is finite.

(24) A is finite iff 2^A is finite.

(25) A is finite and for every X such that $X \in A$ holds X is finite iff $\bigcup A$ is finite.

(26) If $\operatorname{dom} f$ is finite, then $\operatorname{rng} f$ is finite.

(27) If $Y \subseteq \operatorname{rng} f$ and $f^{-1}(Y)$ is finite, then Y is finite.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[2] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal2.html.

[3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[4] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[5] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.

[6] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[7] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[8] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received April 6, 1989

Published January 2, 2004