JOURNAL OF FORMALIZED MATHEMATICS
Volume 10, Released 1998, Published 2003
Inst. of Computer Science, Univ. of Białystok

Lattice of Substitutions is a Heyting Algebra

Adam Grabowski
University of Białystok

MML Identifier: HEYTING2.

WWW: http://mizar.org/JFM/Vol10/heyting2.html

The articles [11], [6], [14], [15], [3], [16], [9], [2], [7], [13], [4], [5], [17], [8], [10], [12], and [1] provide the notation and terminology for this paper.

1. Preliminaries

We follow the rules: V, C, x denote sets and A, B denote elements of SubstitutionSet(V,C).

Let a, b be sets. Observe that $\{\langle a, b\rangle\}$ is function-like and relation-like.

One can prove the following propositions:

(1) For all non empty sets V, C there exists an element f of $V \dot{\rightarrow} C$ such that $f \neq \emptyset$.

(2) For all sets a, b such that $b \in$ SubstitutionSet(V,C) and $a \in b$ holds a is a finite function.

(3) For every element f of $V \dot{\rightarrow} C$ and for every set g such that $g \subseteq f$ holds $g \in V \dot{\rightarrow} C$.

(4) $V \dot{\rightarrow} C \subseteq 2^{[:V, C:]}$.

(5) If V is finite and C is finite, then $V \dot{\rightarrow} C$ is finite.

Let us observe that there exists a set which is functional, finite, and non empty.

2. Some Properties of Sets of Substitutions

The following four propositions are true:

(6) For every finite element a of $V \dot{\rightarrow} C$ holds $\{a\} \in$ SubstitutionSet(V,C).

(7) If $A \frown B = A$, then for every set a such that $a \in A$ there exists a set b such that $b \in B$ and $b \subseteq a$.

(8) If $\mu(A \frown B) = A$, then for every set a such that $a \in A$ there exists a set b such that $b \in B$ and $b \subseteq a$.

(9) If for every set a such that $a \in A$ there exists a set b such that $b \in B$ and $b \subseteq a$, then $\mu(A \frown B) = A$.

Let V be a set, let C be a finite set, and let A be an element of Fin$(V \dot{\rightarrow} C)$. The functor Involved$A$ is defined by:

(Def. 1) $x \in$ Involved$A$ iff there exists a finite function f such that $f \in A$ and $x \in \mathrm{dom}\, f$.

In the sequel C is a finite set.

One can prove the following propositions:

(10) For every set V and for every finite set C and for every element A of $\mathrm{Fin}(V \dot{\rightarrow} C)$ holds $\mathrm{Involved} A \subseteq V$.

(11) For every set V and for every finite set C and for every element A of $\mathrm{Fin}(V \dot{\rightarrow} C)$ such that $A = \emptyset$ holds $\mathrm{Involved} A = \emptyset$.

(12) For every set V and for every finite set C and for every element A of $\mathrm{Fin}(V \dot{\rightarrow} C)$ holds $\mathrm{Involved} A$ is finite.

(13) For every finite set C and for every element A of $\mathrm{Fin}(\emptyset \dot{\rightarrow} C)$ holds $\mathrm{Involved} A = \emptyset$.

Let V be a set, let C be a finite set, and let A be an element of $\mathrm{Fin}(V \dot{\rightarrow} C)$. The functor $-A$ yields an element of $\mathrm{Fin}(V \dot{\rightarrow} C)$ and is defined as follows:

(Def. 2) $-A = \{f; f$ ranges over elements of $\mathrm{Involved} A \dot{\rightarrow} C : \bigwedge_{g:\text{element of } V \dot{\rightarrow} C} (g \in A \Rightarrow f \not\approx g)\}$.

One can prove the following propositions:

(14) $A \frown -A = \emptyset$.

(15) If $A = \emptyset$, then $-A = \{\emptyset\}$.

(16) If $A = \{\emptyset\}$, then $-A = \emptyset$.

(17) For every set V and for every finite set C and for every element A of $\mathrm{SubstitutionSet}(V, C)$ holds $\mu(A \frown -A) = \bot_{\mathrm{SubstLatt}(V,C)}$.

(18) For every non empty set V and for every finite non empty set C and for every element A of $\mathrm{SubstitutionSet}(V, C)$ such that $A = \emptyset$ holds $\mu(-A) = \top_{\mathrm{SubstLatt}(V,C)}$.

(19) Let V be a set, C be a finite set, A be an element of $\mathrm{SubstitutionSet}(V, C)$, a be an element of $V \dot{\rightarrow} C$, and B be an element of $\mathrm{SubstitutionSet}(V, C)$. Suppose $B = \{a\}$. If $A \frown B = \emptyset$, then there exists a finite set b such that $b \in -A$ and $b \subseteq a$.

Let V be a set, let C be a finite set, and let A, B be elements of $\mathrm{Fin}(V \dot{\rightarrow} C)$. The functor $A \rightarrowtail B$ yields an element of $\mathrm{Fin}(V \dot{\rightarrow} C)$ and is defined as follows:

(Def. 3) $A \rightarrowtail B = (V \dot{\rightarrow} C) \cap \{\bigcup\{f(i) \setminus i; i$ ranges over elements of $V \dot{\rightarrow} C : i \in A\}; f$ ranges over elements of $A \dot{\rightarrow} B : \mathrm{dom} f = A\}$.

The following two propositions are true:

(20) Let A, B be elements of $\mathrm{Fin}(V \dot{\rightarrow} C)$ and s be a set. Suppose $s \in A \rightarrowtail B$. Then there exists a partial function f from A to B such that $s = \bigcup\{f(i) \setminus i; i$ ranges over elements of $V \dot{\rightarrow} C : i \in A\}$ and $\mathrm{dom} f = A$.

(21) For every set V and for every finite set C and for every element A of $\mathrm{Fin}(V \dot{\rightarrow} C)$ such that $A = \emptyset$ holds $A \rightarrowtail A = \{\emptyset\}$.

We adopt the following rules: u, v are elements of $\mathrm{SubstLatt}(V, C)$, a is an element of $V \dot{\rightarrow} C$, and K, L are elements of $\mathrm{SubstitutionSet}(V, C)$.

One can prove the following proposition

(22) For every set X such that $X \subseteq u$ holds X is an element of $\mathrm{SubstLatt}(V, C)$.

3. Lattice of Substitutions is Implicative

Let us consider V, C. The functor pseudo_compl(V,C) yields a unary operation on the carrier of SubstLatt(V,C) and is defined as follows:

(Def. 4) For every element u' of SubstitutionSet(V,C) such that $u' = u$ holds $(\text{pseudo\_compl}(V,C))(u) = \mu(-u')$.

The functor StrongImpl(V,C) yields a binary operation on the carrier of SubstLatt(V,C) and is defined by:

(Def. 5) For all elements u', v' of SubstitutionSet(V,C) such that $u' = u$ and $v' = v$ holds $(\text{StrongImpl}(V,C))(u, v) = \mu(u' \rightarrowtail v')$.

Let us consider u. The functor 2^u yielding an element of Fin(the carrier of SubstLatt(V,C)) is defined by:

(Def. 6) $2^u = 2^u$.

The functor $\Box \setminus_u \Box$ yielding a unary operation on the carrier of SubstLatt(V,C) is defined as follows:

(Def. 7) $(\Box \setminus_u \Box)(v) = u \setminus v$.

Let us consider V, C. The functor Atom(V,C) yielding a function from $V \dot{\rightarrow} C$ into the carrier of SubstLatt(V,C) is defined by:

(Def. 8) For every element a of $V \dot{\rightarrow} C$ holds $(\text{Atom}(V,C))(a) = \mu\{a\}$.

One can prove the following propositions:

(23) $\bigsqcup^{\text{f}}_{K} \text{Atom}(V,C) = \text{FinUnion}(K, \text{singleton}_{V \dot{\rightarrow} C})$.

(24) For every element u of SubstitutionSet(V,C) holds $u = \bigsqcup^{\text{f}}_{u} \text{Atom}(V,C)$.

(25) $(\Box \setminus_u \Box)(v) \sqsubseteq u$.

(26) For every element a of $V \dot{\rightarrow} C$ such that a is finite and for every set c such that $c \in (\text{Atom}(V,C))(a)$ holds $c = a$.

(27) For every element a of $V \dot{\rightarrow} C$ such that $K = \{a\}$ and $L = u$ and $L \frown K = \emptyset$ holds $(\text{Atom}(V,C))(a) \sqsubseteq (\text{pseudo\_compl}(V,C))(u)$.

(28) For every finite element a of $V \dot{\rightarrow} C$ holds $a \in (\text{Atom}(V,C))(a)$.

(29) Let u, v be elements of SubstitutionSet(V,C). Suppose that for every set c such that $c \in u$ there exists a set b such that $b \in v$ and $b \subseteq c \cup a$. Then there exists a set b such that $b \in u \rightarrowtail v$ and $b \subseteq a$.

(30) Let a be a finite element of $V \dot{\rightarrow} C$. Suppose for every element b of $V \dot{\rightarrow} C$ such that $b \in u$ holds $b \approx a$ and $u \sqcap (\text{Atom}(V,C))(a) \sqsubseteq v$. Then $(\text{Atom}(V,C))(a) \sqsubseteq (\text{StrongImpl}(V,C))(u, v)$.

(31) $u \sqcap (\text{pseudo\_compl}(V,C))(u) = \bot_{\text{SubstLatt}(V,C)}$.

(32) $u \sqcap (\text{StrongImpl}(V,C))(u, v) \sqsubseteq v$.

Let us consider V, C. Observe that SubstLatt(V,C) is implicative.
Next we state the proposition

(33) $u \Rightarrow v = \bigsqcup^{\text{f}}_{2^u}((\text{the meet operation of SubstLatt}(V,C))^\circ(\text{pseudo\_compl}(V,C), (\text{StrongImpl}(V,C))^\circ(\Box \setminus_u \Box, v)))$.

REFERENCES

[1] Grzegorz Bancerek. Filters — part I. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/filter_0.html.

[2] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.

[3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[4] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[5] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/partfun1.html.

[6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[7] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[8] Adam Grabowski. Lattice of substitutions. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/substlat.html.

[9] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funcop_1.html.

[10] Andrzej Trybulec. Semilattice operations on finite subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setwiseo.html.

[11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[12] Andrzej Trybulec. Finite join and finite meet, and dual lattices. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/lattice2.html.

[13] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finsub_1.html.

[14] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[15] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[16] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

[17] Stanisław Żukowski. Introduction to lattice theory. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/lattices.html.

Received December 31, 1998

Published January 2, 2004