JOURNAL OF FORMALIZED MATHEMATICS
Volume 6, Released 1994, Published 2003
Inst. of Computer Science, Univ. of Białystok

Some Properties of the Intervals

Józef Białas
Łódź University

MML Identifier: MEASURE6.

WWW: http://mizar.org/JFM/Vol6/measure6.html

The articles [8], [7], [10], [1], [9], [11], [5], [6], [2], [3], and [4] provide the notation and terminology for this paper.

The following propositions are true:

(1) There exists a function F from $\mathbb{N}$ into $[:\mathbb{N}, \mathbb{N}:]$ such that F is one-to-one and $\operatorname{dom} F = \mathbb{N}$ and $\operatorname{rng} F = [:\mathbb{N}, \mathbb{N}:]$.

(2) For every function F from $\mathbb{N}$ into $\overline{\mathbb{R}}$ such that F is non-negative holds $0_{\overline{\mathbb{R}}} \leq \sum F$.

(3) Let F be a function from $\mathbb{N}$ into $\overline{\mathbb{R}}$ and x be an extended real number. Suppose there exists a natural number n such that $x \leq F(n)$ and F is non-negative. Then $x \leq \sum F$.

(8)[1] For all extended real numbers x, y such that x is a real number holds $(y-x)+x = y$ and $(y+x)-x = y$.

(10)[2] For all extended real numbers x, y, z such that $z \in \mathbb{R}$ and $y < x$ holds $(z+x)-(z+y) = x-y$.

(11) For all extended real numbers x, y, z such that $z \in \mathbb{R}$ and $x \leq y$ holds $z+x \leq z+y$ and $x+z \leq y+z$ and $x-z \leq y-z$.

(12) For all extended real numbers x, y, z such that $z \in \mathbb{R}$ and $x < y$ holds $z+x < z+y$ and $x+z < y+z$ and $x-z < y-z$.

Let x be a real number. The functor $\overline{\mathbb{R}}(x)$ yielding an extended real number is defined as follows:

(Def. 1) $\overline{\mathbb{R}}(x) = x$.

We now state a number of propositions:

(13) For all real numbers x, y holds $x \leq y$ iff $\overline{\mathbb{R}}(x) \leq \overline{\mathbb{R}}(y)$.

(14) For all real numbers x, y holds $x < y$ iff $\overline{\mathbb{R}}(x) < \overline{\mathbb{R}}(y)$.

(15) For all extended real numbers x, y, z such that $x < y$ and $y < z$ holds y is a real number.

(16) Let x, y, z be extended real numbers. Suppose x is a real number and z is a real number and $x \leq y$ and $y \leq z$. Then y is a real number.

(17) For all extended real numbers x, y, z such that x is a real number and $x \leq y$ and $y < z$ holds y is a real number.

[1] The propositions (4)–(7) have been removed.
[2] The proposition (9) has been removed.

(18) For all extended real numbers x, y, z such that $x < y$ and $y \leq z$ and z is a real number holds y is a real number.

(19) For all extended real numbers x, y such that $0_{\overline{\mathbb{R}}} < x$ and $x < y$ holds $0_{\overline{\mathbb{R}}} < y - x$.

(20) For all extended real numbers x, y, z such that $0_{\overline{\mathbb{R}}} \leq x$ and $0_{\overline{\mathbb{R}}} \leq z$ and $z + x < y$ holds $z < y - x$.

(21) For every extended real number x holds $x - 0_{\overline{\mathbb{R}}} = x$.

(22) For all extended real numbers x, y, z such that $0_{\overline{\mathbb{R}}} \leq x$ and $0_{\overline{\mathbb{R}}} \leq z$ and $z + x < y$ holds $z \leq y$.

(23) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ there exists an extended real number y such that $0_{\overline{\mathbb{R}}} < y$ and $y < x$.

(24) Let x, z be extended real numbers. Suppose $0_{\overline{\mathbb{R}}} < x$ and $x < z$. Then there exists an extended real number y such that $0_{\overline{\mathbb{R}}} < y$ and $x + y < z$ and $y \in \mathbb{R}$.

(25) Let x, z be extended real numbers. Suppose $0_{\overline{\mathbb{R}}} \leq x$ and $x < z$. Then there exists an extended real number y such that $0_{\overline{\mathbb{R}}} < y$ and $x + y < z$ and $y \in \mathbb{R}$.

(26) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ there exists an extended real number y such that $0_{\overline{\mathbb{R}}} < y$ and $y + y < x$.

Let x be an extended real number. Let us assume that $0_{\overline{\mathbb{R}}} < x$. The functor $\mathrm{Seg}\,x$ yields a non empty subset of $\overline{\mathbb{R}}$ and is defined by:

(Def. 2) For every extended real number y holds $y \in \mathrm{Seg}\,x$ iff $0_{\overline{\mathbb{R}}} < y$ and $y + y < x$.

Let x be an extended real number. The functor $\mathrm{len}\,x$ yields an extended real number and is defined by:

(Def. 3) $\mathrm{len}\,x = \sup \mathrm{Seg}\,x$.

Next we state several propositions:

(27) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ holds $0_{\overline{\mathbb{R}}} < \mathrm{len}\,x$.

(28) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ holds $\mathrm{len}\,x \leq x$.

(29) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ and $x < +\infty$ holds $\mathrm{len}\,x$ is a real number.

(30) For every extended real number x such that $0_{\overline{\mathbb{R}}} < x$ holds $\mathrm{len}\,x + \mathrm{len}\,x \leq x$.

(31) Let e_1 be an extended real number. Suppose $0_{\overline{\mathbb{R}}} < e_1$. Then there exists a function F from $\mathbb{N}$ into $\overline{\mathbb{R}}$ such that for every natural number n holds $0_{\overline{\mathbb{R}}} < F(n)$ and $\sum F < e_1$.

(32) Let e_1 be an extended real number and X be a non empty subset of $\overline{\mathbb{R}}$. Suppose $0_{\overline{\mathbb{R}}} < e_1$ and $\inf X$ is a real number. Then there exists an extended real number x such that $x \in X$ and $x < \inf X + e_1$.

(33) Let e_1 be an extended real number and X be a non empty subset of $\overline{\mathbb{R}}$. Suppose $0_{\overline{\mathbb{R}}} < e_1$ and $\sup X$ is a real number. Then there exists an extended real number x such that $x \in X$ and $\sup X - e_1 < x$.

(34) Let F be a function from $\mathbb{N}$ into $\overline{\mathbb{R}}$. Suppose F is non-negative and $\sum F < +\infty$. Let n be a natural number. Then $F(n) \in \mathbb{R}$.

$-\infty$ is an extended real number. Then $+\infty$ is an extended real number.
One can prove the following propositions:

(35) $\mathbb{R}$ is an interval and $\mathbb{R} =]-\infty, +\infty[$ and $\mathbb{R} = [-\infty, +\infty]$ and $\mathbb{R} = [-\infty, +\infty[$ and $\mathbb{R} =]-\infty, +\infty]$.

(36) For all extended real numbers a, b such that $b = -\infty$ holds $]a,b[= \emptyset$ and $[a,b] = \emptyset$ and $[a,b[= \emptyset$ and $]a,b] = \emptyset$.

(37) For all extended real numbers a, b such that $a = +\infty$ holds $]a,b[= \emptyset$ and $[a,b] = \emptyset$ and $[a,b[= \emptyset$ and $]a,b] = \emptyset$.

(38) Let A be an interval, a, b be extended real numbers, and c, d, e be real numbers. If $A =]a,b[$ and $c \in A$ and $d \in A$ and $c \leq e$ and $e \leq d$, then $e \in A$.

(39) Let A be an interval, a, b be extended real numbers, and c, d, e be real numbers. If $A = [a,b]$ and $c \in A$ and $d \in A$ and $c \leq e$ and $e \leq d$, then $e \in A$.

(40) Let A be an interval, a, b be extended real numbers, and c, d, e be real numbers. If $A =]a,b]$ and $c \in A$ and $d \in A$ and $c \leq e$ and $e \leq d$, then $e \in A$.

(41) Let A be an interval, a, b be extended real numbers, and c, d, e be real numbers. If $A = [a,b[$ and $c \in A$ and $d \in A$ and $c \leq e$ and $e \leq d$, then $e \in A$.

(42) Let A be a non empty subset of $\overline{\mathbb{R}}$ and m, M be extended real numbers. Suppose $m = \inf A$ and $M = \sup A$. Suppose that

(i) for all real numbers c, d such that $c \in A$ and $d \in A$ and for every real number e such that $c \leq e$ and $e \leq d$ holds $e \in A$,

(ii) $m \notin A$, and

(iii) $M \notin A$.

Then $A =]m,M[$.

(43) Let A be a non empty subset of $\overline{\mathbb{R}}$ and m, M be extended real numbers. Suppose $m = \inf A$ and $M = \sup A$. Suppose that

(i) for all real numbers c, d such that $c \in A$ and $d \in A$ and for every real number e such that $c \leq e$ and $e \leq d$ holds $e \in A$,

(ii) $m \in A$,

(iii) $M \in A$, and

(iv) $A \subseteq \mathbb{R}$.

Then $A = [m,M]$.

(44) Let A be a non empty subset of $\overline{\mathbb{R}}$ and m, M be extended real numbers. Suppose $m = \inf A$ and $M = \sup A$. Suppose that

(i) for all real numbers c, d such that $c \in A$ and $d \in A$ and for every real number e such that $c \leq e$ and $e \leq d$ holds $e \in A$,

(ii) $m \in A$,

(iii) $M \notin A$, and

(iv) $A \subseteq \mathbb{R}$.

Then $A = [m,M[$.

(45) Let A be a non empty subset of $\overline{\mathbb{R}}$ and m, M be extended real numbers. Suppose $m = \inf A$ and $M = \sup A$. Suppose that

(i) for all real numbers c, d such that $c \in A$ and $d \in A$ and for every real number e such that $c \leq e$ and $e \leq d$ holds $e \in A$,

(ii) $m \notin A$,

(iii) $M \in A$, and

(iv) $A \subseteq \mathbb{R}$.

Then $A =]m,M]$.

(46) Let A be a subset of $\mathbb{R}$. Then A is an interval if and only if for all real numbers a, b such that $a \in A$ and $b \in A$ and for every real number c such that $a \leq c$ and $c \leq b$ holds $c \in A$.

(47) For all intervals A, B such that A meets B holds $A \cup B$ is an interval.

Let A be an interval. Let us assume that $A \neq \emptyset$. The functor $\inf A$ yields an extended real number and is defined as follows:

(Def. 4) There exists an extended real number b such that $\inf A \leq b$ but $A =]\inf A, b[$ or $A =]\inf A, b]$ or $A = [\inf A, b]$ or $A = [\inf A, b[$.

Let A be an interval. Let us assume that $A \neq \emptyset$. The functor $\sup A$ yields an extended real number and is defined as follows:

(Def. 5) There exists an extended real number a such that $a \leq \sup A$ but $A =]a, \sup A[$ or $A =]a, \sup A]$ or $A = [a, \sup A]$ or $A = [a, \sup A[$.

One can prove the following propositions:

(48) For every interval A such that A is open interval and $A \neq \emptyset$ holds $\inf A \leq \sup A$ and $A =]\inf A, \sup A[$.

(49) For every interval A such that A is closed interval and $A \neq \emptyset$ holds $\inf A \leq \sup A$ and $A = [\inf A, \sup A]$.

(50) For every interval A such that A is right open interval and $A \neq \emptyset$ holds $\inf A \leq \sup A$ and $A = [\inf A, \sup A[$.

(51) For every interval A such that A is left open interval and $A \neq \emptyset$ holds $\inf A \leq \sup A$ and $A =]\inf A, \sup A]$.

(52) For every interval A such that $A \neq \emptyset$ holds $\inf A \leq \sup A$ but $A =]\inf A, \sup A[$ or $A =]\inf A, \sup A]$ or $A = [\inf A, \sup A]$ or $A = [\inf A, \sup A[$.

(54)[3] For every interval A and for every real number a such that $a \in A$ holds $\inf A \leq \overline{\mathbb{R}}(a)$ and $\overline{\mathbb{R}}(a) \leq \sup A$.

(55) For all intervals A, B and for all real numbers a, b such that $a \in A$ and $b \in B$ holds if $\sup A \leq \inf B$, then $a \leq b$.

(56) For every interval A and for every extended real number a such that $a \in A$ holds $\inf A \leq a$ and $a \leq \sup A$.

(57) For every interval A such that $A \neq \emptyset$ and for every extended real number a such that $\inf A < a$ and $a < \sup A$ holds $a \in A$.

(58) For all intervals A, B such that $\sup A = \inf B$ but $\sup A \in A$ or $\inf B \in B$ holds $A \cup B$ is an interval.

Let A be a subset of $\mathbb{R}$ and let x be a real number. The functor $x + A$ yielding a subset of $\mathbb{R}$ is defined as follows:

(Def. 6) For every real number y holds $y \in x + A$ iff there exists a real number z such that $z \in A$ and $y = x + z$.

We now state several propositions:

(59) For every subset A of $\mathbb{R}$ and for every real number x holds $-x + (x + A) = A$.

(60) For every real number x and for every subset A of $\mathbb{R}$ such that $A = \mathbb{R}$ holds $x + A = A$.

(61) For every real number x holds $x + \emptyset = \emptyset$.

[3] The proposition (53) has been removed.

(62) For every interval A and for every real number x holds A is open interval iff $x+A$ is open interval.

(63) For every interval A and for every real number x holds A is closed interval iff $x+A$ is closed interval.

(64) Let A be an interval and x be a real number. Then A is right open interval if and only if $x+A$ is right open interval.

(65) Let A be an interval and x be a real number. Then A is left open interval if and only if $x+A$ is left open interval.

(66) For every interval A and for every real number x holds $x+A$ is an interval.

Let A be an interval and let x be a real number. One can check that $x+A$ is interval.
We now state the proposition

(67) For every interval A and for every real number x holds $\mathrm{vol}(A) = \mathrm{vol}(x+A)$.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[2] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/supinf_1.html.

[3] Józef Białas. Series of positive real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/supinf_2.html.

[4] Józef Białas. Properties of the intervals of real numbers. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/measure5.html.

[5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[8] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[9] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[10] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[11] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received February 5, 1994

Published January 2, 2004