JOURNAL OF FORMALIZED MATHEMATICS
Volume 5, Released 1993, Published 2003
Inst. of Computer Science, Univ. of Białystok

Remarks on Special Subsets of Topological Spaces

Zbigniew Karno
Warsaw University
Białystok

Summary. Let X be a topological space and let A be a subset of X. Recall that A is *nowhere dense* in X if its closure is a boundary subset of X, i.e., if $\mathrm{Int}\overline{A} = \emptyset$ (see [2]). We introduce here the concept of everywhere dense subsets in X, which is dual to the above one. Namely, A is said to be *everywhere dense* in X if its interior is a dense subset of X, i.e., if $\overline{\mathrm{Int}A} =$ the carrier of X.

Our purpose is to list a number of properties of such sets (comp. [7]). As a sample we formulate their two dual characterizations. The first one characterizes thin sets in X : *A is nowhere dense iff for every open nonempty subset G of X there is an open nonempty subset of X contained in G and disjoint from A*. The corresponding second one characterizes thick sets in X : *A is everywhere dense iff for every closed subset F of X distinct from the carrier of X there is a closed subset of X distinct from the carrier of X, which contains F and together with A covers the carrier of X*. We also give some connections between both these concepts. Of course, *A is everywhere (nowhere) dense in X iff its complement is nowhere (everywhere) dense*. Moreover, *A is nowhere dense iff there are two subsets of X, C boundary closed and B everywhere dense, such that $A = C \cap B$ and $C \cup B$ covers the carrier of X*. Dually, *A is everywhere dense iff there are two disjoint subsets of X, C open dense and B nowhere dense, such that $A = C \cup B$*.

Note that some relationships between everywhere (nowhere) dense sets in X and everywhere (nowhere) dense sets in subspaces of X are also indicated.

MML Identifier: `TOPS_3`.

WWW: `http://mizar.org/JFM/Vol5/tops_3.html`

The articles [4], [6], [3], [7], [5], and [1] provide the notation and terminology for this paper.

1. Selected Properties of Subsets of a Topological Space

In this paper X denotes a topological structure and A denotes a subset of X.

We now state three propositions:

(1) $A = \emptyset_X$ iff $A^{\rm c} = \Omega_X$ and $A = \emptyset$ iff $A^{\rm c} =$ the carrier of X.

(2) $A = \Omega_X$ iff $A^{\rm c} = \emptyset_X$ and $A =$ the carrier of X iff $A^{\rm c} = \emptyset$.

(3) For every topological space X and for all subsets A, B of X holds $\mathrm{Int}A \cap \overline{B} \subseteq \overline{A \cap B}$.

In the sequel X denotes a topological space and A, B denote subsets of X.

We now state several propositions:

(4) $\mathrm{Int}(A \cup B) \subseteq \overline{A} \cup \mathrm{Int}B$.

(5) For every subset A of X such that A is closed holds $\mathrm{Int}(A \cup B) \subseteq A \cup \mathrm{Int}B$.

(6) For every subset A of X such that A is closed holds $\mathrm{Int}(A \cup B) = \mathrm{Int}(A \cup \mathrm{Int} B)$.

(7) If A misses $\mathrm{Int}\overline{A}$, then $\mathrm{Int}\overline{A} = \emptyset$.

(8) If $A \cup \overline{\mathrm{Int} A} =$ the carrier of X, then $\overline{\mathrm{Int} A} =$ the carrier of X.

2. Special Subsets of a Topological Space

Let X be a topological structure and let A be a subset of X. Let us observe that A is boundary if and only if:

(Def. 1) $\mathrm{Int} A = \emptyset$.

We now state the proposition

(9) $\emptyset_X$ is boundary.

In the sequel X is a non empty topological space and A is a subset of X.
Next we state the proposition

(10) If A is boundary, then $A \neq$ the carrier of X.

In the sequel X is a topological space and A, B are subsets of X.
We now state several propositions:

(11) If B is boundary and $A \subseteq B$, then A is boundary.

(12) A is boundary iff for every subset C of X such that $A^{\mathrm{c}} \subseteq C$ and C is closed holds $C =$ the carrier of X.

(13) A is boundary iff for every subset G of X such that $G \neq \emptyset$ and G is open holds A^{c} meets G.

(14) A is boundary iff for every subset F of X such that F is closed holds $\mathrm{Int} F = \mathrm{Int}(F \cup A)$.

(15) If A is boundary or B is boundary, then $A \cap B$ is boundary.

Let X be a topological structure and let A be a subset of X. Let us observe that A is dense if and only if:

(Def. 2) $\overline{A} =$ the carrier of X.

Next we state the proposition

(16) Ω_X is dense.

In the sequel X denotes a non empty topological space and A, B denote subsets of X.
The following propositions are true:

(17) If A is dense, then $A \neq \emptyset$.

(18) A is dense iff A^{c} is boundary.

(19) A is dense iff for every subset C of X such that $A \subseteq C$ and C is closed holds $C =$ the carrier of X.

(20) A is dense iff for every subset G of X such that G is open holds $\overline{G} = \overline{G \cap A}$.

(21) If A is dense or B is dense, then $A \cup B$ is dense.

Let X be a topological structure and let A be a subset of X. Let us observe that A is nowhere dense if and only if:

(Def. 3) $\mathrm{Int}\overline{A} = \emptyset$.

The following propositions are true:

(22) $\emptyset_X$ is nowhere dense.

(23) If A is nowhere dense, then $A \neq$ the carrier of X.

(24) If A is nowhere dense, then $\overline{A}$ is nowhere dense.

(25) If A is nowhere dense, then A is not dense.

(26) If B is nowhere dense and $A \subseteq B$, then A is nowhere dense.

(27) A is nowhere dense iff there exists a subset C of X such that $A \subseteq C$ and C is closed and boundary.

(28) A is nowhere dense if and only if for every subset G of X such that $G \neq \emptyset$ and G is open there exists a subset H of X such that $H \subseteq G$ and $H \neq \emptyset$ and H is open and A misses H.

(29) If A is nowhere dense or B is nowhere dense, then $A \cap B$ is nowhere dense.

(30) If A is nowhere dense and B is boundary, then $A \cup B$ is boundary.

Let X be a topological structure and let A be a subset of X. We say that A is everywhere dense if and only if:

(Def. 4) $\overline{\mathrm{Int} A} = \Omega_X$.

Let X be a topological structure and let A be a subset of X. Let us observe that A is everywhere dense if and only if:

(Def. 5) $\overline{\mathrm{Int} A} =$ the carrier of X.

We now state a number of propositions:

(31) Ω_X is everywhere dense.

(32) If A is everywhere dense, then $\mathrm{Int} A$ is everywhere dense.

(33) If A is everywhere dense, then A is dense.

(34) If A is everywhere dense, then $A \neq \emptyset$.

(35) A is everywhere dense iff $\mathrm{Int} A$ is dense.

(36) If A is open and dense, then A is everywhere dense.

(37) If A is everywhere dense, then A is not boundary.

(38) If A is everywhere dense and $A \subseteq B$, then B is everywhere dense.

(39) A is everywhere dense iff A^{c} is nowhere dense.

(40) A is nowhere dense iff A^{c} is everywhere dense.

(41) A is everywhere dense iff there exists a subset C of X such that $C \subseteq A$ and C is open and dense.

(42) A is everywhere dense if and only if for every subset F of X such that $F \neq$ the carrier of X and F is closed there exists a subset H of X such that $F \subseteq H$ and $H \neq$ the carrier of X and H is closed and $A \cup H =$ the carrier of X.

(43) If A is everywhere dense or B is everywhere dense, then $A \cup B$ is everywhere dense.

(44) If A is everywhere dense and B is everywhere dense, then $A \cap B$ is everywhere dense.

(45) If A is everywhere dense and B is dense, then $A \cap B$ is dense.

(46) If A is dense and B is nowhere dense, then $A \setminus B$ is dense.

(47) If A is everywhere dense and B is boundary, then $A \setminus B$ is dense.

(48) If A is everywhere dense and B is nowhere dense, then $A \setminus B$ is everywhere dense.

In the sequel D is a subset of X.
The following four propositions are true:

(49) Suppose D is everywhere dense. Then there exist subsets C, B of X such that C is open and dense and B is nowhere dense and $C \cup B = D$ and C misses B.

(50) Suppose D is everywhere dense. Then there exist subsets C, B of X such that C is open and dense and B is closed and boundary and $C \cup D \cap B = D$ and C misses B and $C \cup B =$ the carrier of X.

(51) Suppose D is nowhere dense. Then there exist subsets C, B of X such that C is closed and boundary and B is everywhere dense and $C \cap B = D$ and $C \cup B =$ the carrier of X.

(52) Suppose D is nowhere dense. Then there exist subsets C, B of X such that C is closed and boundary and B is open and dense and $C \cap (D \cup B) = D$ and C misses B and $C \cup B =$ the carrier of X.

3. Properties of Subsets in Subspaces

In the sequel Y_0 is a subspace of X.
The following propositions are true:

(53) For every subset A of X and for every subset B of Y_0 such that $B \subseteq A$ holds $\overline{B} \subseteq \overline{A}$.

(54) Let C, A be subsets of X and B be a subset of Y_0. If C is closed and $C \subseteq$ the carrier of Y_0 and $A \subseteq C$ and $A = B$, then $\overline{A} = \overline{B}$.

(55) Let Y_0 be a closed non empty subspace of X, A be a subset of X, and B be a subset of Y_0. If $A = B$, then $\overline{A} = \overline{B}$.

(56) For every subset A of X and for every subset B of Y_0 such that $A \subseteq B$ holds $\mathrm{Int} A \subseteq \mathrm{Int} B$.

(57) Let Y_0 be a non empty subspace of X, C, A be subsets of X, and B be a subset of Y_0. If C is open and $C \subseteq$ the carrier of Y_0 and $A \subseteq C$ and $A = B$, then $\mathrm{Int} A = \mathrm{Int} B$.

(58) Let Y_0 be an open non empty subspace of X, A be a subset of X, and B be a subset of Y_0. If $A = B$, then $\mathrm{Int} A = \mathrm{Int} B$.

In the sequel X_0 is a subspace of X.
Next we state two propositions:

(59) For every subset A of X and for every subset B of X_0 such that $A \subseteq B$ holds if A is dense, then B is dense.

(60) Let C, A be subsets of X and B be a subset of X_0. Suppose $C \subseteq$ the carrier of X_0 and $A \subseteq C$ and $A = B$. Then C is dense and B is dense if and only if A is dense.

In the sequel X_0 denotes a non empty subspace of X.
Next we state a number of propositions:

(61) Let A be a subset of X and B be a subset of X_0. If $A \subseteq B$, then if A is everywhere dense, then B is everywhere dense.

(62) Let C, A be subsets of X and B be a subset of X_0. Suppose C is open and $C \subseteq$ the carrier of X_0 and $A \subseteq C$ and $A = B$. Then C is dense and B is everywhere dense if and only if A is everywhere dense.

(63) Let X_0 be an open non empty subspace of X, A, C be subsets of X, and B be a subset of X_0. Suppose $C =$ the carrier of X_0 and $A = B$. Then C is dense and B is everywhere dense if and only if A is everywhere dense.

(64) Let C, A be subsets of X and B be a subset of X_0. Suppose $C \subseteq$ the carrier of X_0 and $A \subseteq C$ and $A = B$. Then C is everywhere dense and B is everywhere dense if and only if A is everywhere dense.

(65) For every subset A of X and for every subset B of X_0 such that $A \subseteq B$ holds if B is boundary, then A is boundary.

(66) Let C, A be subsets of X and B be a subset of X_0. Suppose C is open and $C \subseteq$ the carrier of X_0 and $A \subseteq C$ and $A = B$. If A is boundary, then B is boundary.

(67) Let X_0 be an open non empty subspace of X, A be a subset of X, and B be a subset of X_0. If $A = B$, then A is boundary iff B is boundary.

(68) Let A be a subset of X and B be a subset of X_0. If $A \subseteq B$, then if B is nowhere dense, then A is nowhere dense.

(69) Let C, A be subsets of X and B be a subset of X_0. Suppose C is open and $C \subseteq$ the carrier of X_0 and $A \subseteq C$ and $A = B$. If A is nowhere dense, then B is nowhere dense.

(70) Let X_0 be an open non empty subspace of X, A be a subset of X, and B be a subset of X_0. If $A = B$, then A is nowhere dense iff B is nowhere dense.

4. Subsets in Topological Spaces with the same Topological Structures

We now state the proposition

(71) Let X_1, X_2 be 1-sorted structures. Suppose the carrier of $X_1 =$ the carrier of X_2. Let C_1 be a subset of X_1 and C_2 be a subset of X_2. Then $C_1 = C_2$ if and only if ${C_1}^{\mathrm{c}} = {C_2}^{\mathrm{c}}$.

In the sequel X_1, X_2 are topological structures.

Next we state two propositions:

(72) Suppose that

(i) the carrier of $X_1 =$ the carrier of X_2, and

(ii) for every subset C_1 of X_1 and for every subset C_2 of X_2 such that $C_1 = C_2$ holds C_1 is open iff C_2 is open.

Then the topological structure of $X_1 =$ the topological structure of X_2.

(73) Suppose that

(i) the carrier of $X_1 =$ the carrier of X_2, and

(ii) for every subset C_1 of X_1 and for every subset C_2 of X_2 such that $C_1 = C_2$ holds C_1 is closed iff C_2 is closed.

Then the topological structure of $X_1 =$ the topological structure of X_2.

In the sequel X_1, X_2 are topological spaces.

One can prove the following propositions:

(74) Suppose that

(i) the carrier of $X_1 =$ the carrier of X_2, and

(ii) for every subset C_1 of X_1 and for every subset C_2 of X_2 such that $C_1 = C_2$ holds $\mathrm{Int}\, C_1 = \mathrm{Int}\, C_2$.

Then the topological structure of $X_1 =$ the topological structure of X_2.

(75) Suppose that

(i) the carrier of X_1 = the carrier of X_2, and

(ii) for every subset C_1 of X_1 and for every subset C_2 of X_2 such that $C_1 = C_2$ holds $\overline{C_1} = \overline{C_2}$.

Then the topological structure of X_1 = the topological structure of X_2.

In the sequel D_1 denotes a subset of X_1 and D_2 denotes a subset of X_2.
One can prove the following propositions:

(76) Suppose $D_1 = D_2$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is open, then D_2 is open.

(77) If $D_1 = D_2$ and the topological structure of X_1 = the topological structure of X_2, then $\mathrm{Int}\,D_1 = \mathrm{Int}\,D_2$.

(78) If $D_1 \subseteq D_2$ and the topological structure of X_1 = the topological structure of X_2, then $\mathrm{Int}\,D_1 \subseteq \mathrm{Int}\,D_2$.

(79) Suppose $D_1 = D_2$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is closed, then D_2 is closed.

(80) If $D_1 = D_2$ and the topological structure of X_1 = the topological structure of X_2, then $\overline{D_1} = \overline{D_2}$.

(81) If $D_1 \subseteq D_2$ and the topological structure of X_1 = the topological structure of X_2, then $\overline{D_1} \subseteq \overline{D_2}$.

(82) Suppose $D_2 \subseteq D_1$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is boundary, then D_2 is boundary.

(83) Suppose $D_1 \subseteq D_2$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is dense, then D_2 is dense.

(84) Suppose $D_2 \subseteq D_1$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is nowhere dense, then D_2 is nowhere dense.

In the sequel X_1, X_2 denote non empty topological spaces, D_1 denotes a subset of X_1, and D_2 denotes a subset of X_2.

We now state the proposition

(85) Suppose $D_1 \subseteq D_2$ and the topological structure of X_1 = the topological structure of X_2. If D_1 is everywhere dense, then D_2 is everywhere dense.

REFERENCES

[1] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Journal of Formalized Mathematics*, 4, 1992. `http://mizar.org/JFM/Vol4/tsep_1.html`.

[2] Kazimierz Kuratowski. *Topology*, volume I. PWN - Polish Scientific Publishers, Academic Press, Warsaw, New York and London, 1966.

[3] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/pre_topc.html`.

[4] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[5] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Journal of Formalized Mathematics*, 3, 1991. `http://mizar.org/JFM/Vol3/borsuk_1.html`.

[6] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[7] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_1.html.

Received April 6, 1993

Published January 2, 2004