JOURNAL OF FORMALIZED MATHEMATICS
Volume 13, Released 2001, Published 2003
Inst. of Computer Science, Univ. of Białystok

Some Properties of Dyadic Numbers and Intervals

Józef Białas
Łódź University

Yatsuka Nakamura
Shinshu University
Nagano

Summary. The article is the second part of a paper proving the fundamental Urysohn Theorem concerning the existence of a real valued continuous function on a normal topological space. The paper is divided into two parts. In the first part, we introduce some definitions and theorems concerning properties of intervals; in the second we prove some of properties of dyadic numbers used in proving Urysohn Lemma.

MML Identifier: URYSOHN2.

WWW: http://mizar.org/JFM/Vol13/urysohn2.html

The articles [11], [13], [1], [8], [12], [9], [2], [3], [4], [5], [6], [10], and [7] provide the notation and terminology for this paper.

One can prove the following propositions:

(1) For every interval A such that $A \neq \emptyset$ holds if $\inf A < \sup A$, then $\mathrm{vol}(A) = \sup A - \inf A$ and if $\sup A = \inf A$, then $\mathrm{vol}(A) = 0_{\overline{\mathbb{R}}}$.

(2) For every subset A of $\mathbb{R}$ and for every real number x such that $x \neq 0$ holds $x^{-1} \cdot (x \cdot A) = A$.

(3) For every real number x such that $x \neq 0$ and for every subset A of $\mathbb{R}$ such that $A = \mathbb{R}$ holds $x \cdot A = A$.

(4) For every subset A of $\mathbb{R}$ such that $A \neq \emptyset$ holds $0 \cdot A = \{0\}$.

(5) For every real number x holds $x \cdot \emptyset = \emptyset$.

(6) Let a, b be extended real numbers. Suppose $a \leq b$. Then $a = -\infty$ and $b = -\infty$ or $a = -\infty$ and $b \in \mathbb{R}$ or $a = -\infty$ and $b = +\infty$ or $a \in \mathbb{R}$ and $b \in \mathbb{R}$ or $a \in \mathbb{R}$ and $b = +\infty$ or $a = +\infty$ and $b = +\infty$.

(7) For every extended real number x holds $[x,x]$ is an interval.

(8) For every interval A holds $0 \cdot A$ is an interval.

(9) Let A be an interval and x be a real number. If $x \neq 0$, then if A is open interval, then $x \cdot A$ is open interval.

(10) Let A be an interval and x be a real number. If $x \neq 0$, then if A is closed interval, then $x \cdot A$ is closed interval.

(11) Let A be an interval and x be a real number. Suppose $0 < x$. If A is right open interval, then $x \cdot A$ is right open interval.

(12) Let A be an interval and x be a real number. Suppose $x < 0$. If A is right open interval, then $x \cdot A$ is left open interval.

(13) Let A be an interval and x be a real number. Suppose $0 < x$. If A is left open interval, then $x \cdot A$ is left open interval.

(14) Let A be an interval and x be a real number. Suppose $x < 0$. If A is left open interval, then $x \cdot A$ is right open interval.

(15) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A]$. Then $B = [\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.

(16) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A]$. Then $B =]\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.

(17) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A[$. Then $B =]\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.

(18) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A[$. Then $B = [\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.

(19) For every interval A and for every real number x holds $x \cdot A$ is an interval.

Let A be an interval and let x be a real number. Note that $x \cdot A$ is interval.
One can prove the following propositions:

(20) Let A be an interval and x be a real number. If $0 \leq x$, then for every real number y such that $y = \mathrm{vol}(A)$ holds $x \cdot y = \mathrm{vol}(x \cdot A)$.

(23)[1] For every real number e_1 such that $0 < e_1$ there exists a natural number n such that $1 < 2^n \cdot e_1$.

(24) For all real numbers a, b such that $0 \leq a$ and $1 < b - a$ there exists a natural number n such that $a < n$ and $n < b$.

(27)[2] For every natural number n holds $\mathrm{dyadic}(n) \subseteq \mathrm{DYADIC}$.

(28) For all real numbers a, b such that $a < b$ and $0 \leq a$ and $b \leq 1$ there exists a real number c such that $c \in \mathrm{DYADIC}$ and $a < c$ and $c < b$.

(29) For all real numbers a, b such that $a < b$ there exists a real number c such that $c \in \mathrm{DOM}$ and $a < c$ and $c < b$.

(30) For every non empty subset A of $\overline{\mathbb{R}}$ and for all extended real numbers a, b such that $A \subseteq [a, b]$ holds $a \leq \inf A$ and $\sup A \leq b$.

(31) $0 \in \mathrm{DYADIC}$ and $1 \in \mathrm{DYADIC}$.

(32) For all extended real numbers a, b such that $a = 0$ and $b = 1$ holds $\mathrm{DYADIC} \subseteq [a, b]$.

(33) For all natural numbers n, k such that $n \leq k$ holds $\mathrm{dyadic}(n) \subseteq \mathrm{dyadic}(k)$.

(34) For all real numbers a, b, c, d such that $a < c$ and $c < b$ and $a < d$ and $d < b$ holds $|d - c| < b - a$.

(35) Let e_1 be a real number. Suppose $0 < e_1$. Let d be a real number. Suppose $0 < d$ and $d \leq 1$. Then there exist real numbers r_1, r_2 such that $r_1 \in \mathrm{DYADIC} \cup \mathbb{R}_{>1}$ and $r_2 \in \mathrm{DYADIC} \cup \mathbb{R}_{>1}$ and $0 < r_1$ and $r_1 < d$ and $d < r_2$ and $r_2 - r_1 < e_1$.

[1] The propositions (21) and (22) have been removed.
[2] The propositions (25) and (26) have been removed.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/ordinal1.html`.

[2] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/supinf_1.html`.

[3] Józef Białas. Series of positive real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/supinf_2.html`.

[4] Józef Białas. Properties of the intervals of real numbers. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/measure5.html`.

[5] Józef Białas. Some properties of the intervals. *Journal of Formalized Mathematics*, 6, 1994. `http://mizar.org/JFM/Vol6/measure6.html`.

[6] Józef Białas and Yatsuka Nakamura. Dyadic numbers and T_4 topological spaces. *Journal of Formalized Mathematics*, 7, 1995. `http://mizar.org/JFM/Vol7/urysohn1.html`.

[7] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Scalar multiple of Riemann definite integral. *Journal of Formalized Mathematics*, 11, 1999. `http://mizar.org/JFM/Vol11/integra2.html`.

[8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/real_1.html`.

[9] Rafał Kwiatek. Factorial and Newton coefficients. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/newton.html`.

[10] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/absvalue.html`.

[11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[12] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. `http://mizar.org/JFM/Addenda/numbers.html`.

[13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

Received February 16, 2001

Published January 2, 2004