JOURNAL OF FORMALIZED MATHEMATICS
Volume 12, Released 2000, Published 2003
Inst. of Computer Science, Univ. of Białystok

The Tichonov Theorem

Bartłomiej Skorulski
University of Białystok

MML Identifier: YELLOW17.

WWW: http://mizar.org/JFM/Vol12/yellow17.html

The articles [15], [5], [17], [18], [4], [3], [8], [16], [12], [9], [1], [13], [7], [6], [14], [10], [2], and [11] provide the notation and terminology for this paper.

1. Some Properties of Products

The following propositions are true:

(1) For every function F and for all sets i, x_1 and for every subset A_1 of $F(i)$ such that $(\mathrm{proj}(F,i))^{-1}(\{x_1\})$ meets $(\mathrm{proj}(F,i))^{-1}(A_1)$ holds $x_1 \in A_1$.

(2) For all functions F, f and for all sets i, x_1 such that $x_1 \in F(i)$ and $f \in \prod F$ holds $f +\cdot (i,x_1) \in \prod F$.

(3) For every function F and for every set i such that $i \in \mathrm{dom} F$ and $\prod F \neq \emptyset$ holds $\mathrm{rng}\,\mathrm{proj}(F,i) = F(i)$.

(4) For every function F and for every set i such that $i \in \mathrm{dom} F$ holds $(\mathrm{proj}(F,i))^{-1}(F(i)) = \prod F$.

(5) For all functions F, f and for all sets i, x_1 such that $x_1 \in F(i)$ and $i \in \mathrm{dom} F$ and $f \in \prod F$ holds $f +\cdot (i,x_1) \in (\mathrm{proj}(F,i))^{-1}(\{x_1\})$.

(6) Let F, f be functions, i_1, i_2, x_2 be sets, and A_2 be a subset of $F(i_2)$. Suppose $x_2 \in F(i_1)$ and $i_1 \in \mathrm{dom} F$ and $f \in \prod F$. If $i_1 \neq i_2$, then $f \in (\mathrm{proj}(F,i_2))^{-1}(A_2)$ iff $f +\cdot (i_1,x_2) \in (\mathrm{proj}(F,i_2))^{-1}(A_2)$.

(7) Let F be a function, i_1, i_2, x_2 be sets, and A_2 be a subset of $F(i_2)$. Suppose $\prod F \neq \emptyset$ and $x_2 \in F(i_1)$ and $i_1 \in \mathrm{dom} F$ and $i_2 \in \mathrm{dom} F$ and $A_2 \neq F(i_2)$. Then $(\mathrm{proj}(F,i_1))^{-1}(\{x_2\}) \subseteq (\mathrm{proj}(F,i_2))^{-1}(A_2)$ if and only if $i_1 = i_2$ and $x_2 \in A_2$.

The scheme *ElProductEx* deals with a non empty set $\mathcal{A}$, a topological space yielding nonempty many sorted set $\mathcal{B}$ indexed by $\mathcal{A}$, and a binary predicate $\mathcal{P}$, and states that:

There exists an element f of $\prod \mathcal{B}$ such that for every element i of $\mathcal{A}$ holds $\mathcal{P}[f(i),i]$

provided the parameters satisfy the following condition:

- For every element i of $\mathcal{A}$ there exists an element x of $\mathcal{B}(i)$ such that $\mathcal{P}[x,i]$.

One can prove the following propositions:

(8) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, and f be an element of $\prod J$. Then $(\mathrm{proj}(J,i))(f) = f(i)$.

(9) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, x_1 be an element of $J(i)$, and A_1 be a subset of $J(i)$. If $(\mathrm{proj}(J,i))^{-1}(\{x_1\})$ meets $(\mathrm{proj}(J,i))^{-1}(A_1)$, then $x_1 \in A_1$.

(10) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, and i be an element of I. Then $(\mathrm{proj}(J,i))^{-1}(\Omega_{J(i)}) = \Omega_{\prod J}$.

(11) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, x_1 be an element of $J(i)$, and f be an element of $\prod J$. Then $f +\cdot (i,x_1) \in (\mathrm{proj}(J,i))^{-1}(\{x_1\})$.

(12) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i_1, i_2 be elements of I, x_2 be an element of $J(i_1)$, and A_2 be a subset of $J(i_2)$. If $A_2 \neq \Omega_{J(i_2)}$, then $(\mathrm{proj}(J,i_1))^{-1}(\{x_2\}) \subseteq (\mathrm{proj}(J,i_2))^{-1}(A_2)$ iff $i_1 = i_2$ and $x_2 \in A_2$.

(13) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i_1, i_2 be elements of I, x_2 be an element of $J(i_1)$, A_2 be a subset of $J(i_2)$, and f be an element of $\prod J$. If $i_1 \neq i_2$, then $f \in (\mathrm{proj}(J,i_2))^{-1}(A_2)$ iff $f +\cdot (i_1,x_2) \in (\mathrm{proj}(J,i_2))^{-1}(A_2)$.

2. Some Properties of Compact Spaces

One can prove the following two propositions:

(15)[1] Let T be a non empty topological structure. Then T is compact if and only if for every family F of subsets of T such that F is open and $\Omega_T \subseteq \bigcup F$ there exists a family G of subsets of T such that $G \subseteq F$ and $\Omega_T \subseteq \bigcup G$ and G is finite.

(16) Let T be a non empty topological space and B be a prebasis of T. Then T is compact if and only if for every subset F of B such that $\Omega_T \subseteq \bigcup F$ there exists a finite subset G of F such that $\Omega_T \subseteq \bigcup G$.

3. The Tichonov Theorem

Next we state several propositions:

(17) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, and A be a set. Suppose $A \in$ the product prebasis for J. Then there exists an element i of I and there exists a subset A_1 of $J(i)$ such that A_1 is open and $(\mathrm{proj}(J,i))^{-1}(A_1) = A$.

(18) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, x_1 be an element of $J(i)$, and A be a set. Suppose $A \in$ the product prebasis for J and $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \subseteq A$. Then $A = \Omega_{\prod J}$ or there exists a subset A_1 of $J(i)$ such that $A_1 \neq \Omega_{J(i)}$ and $x_1 \in A_1$ and A_1 is open and $A = (\mathrm{proj}(J,i))^{-1}(A_1)$.

(19) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, and F_1 be a non empty family of subsets of $J(i)$. If $\Omega_{J(i)} \subseteq \bigcup F_1$, then $\Omega_{\prod J} \subseteq \bigcup \{(\mathrm{proj}(J,i))^{-1}(A_1) : A_1$ ranges over elements of $F_1\}$.

(20) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, x_1 be an element of $J(i)$, and G be a subset of the product prebasis for J. Suppose $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \subseteq \bigcup G$ and for every set A such that $A \in$ the product prebasis for J and $A \in G$ holds $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \not\subseteq A$. Then $\Omega_{\prod J} \subseteq \bigcup G$.

(21) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, and F be a subset of the product prebasis for J. Suppose that for every finite subset G of F holds $\Omega_{\prod J} \not\subseteq \bigcup G$. Let x_1 be an element of $J(i)$ and G be a finite subset of F. Suppose $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \subseteq \bigcup G$. Then there exists a set A such that $A \in$ the product prebasis for J and $A \in G$ and $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \subseteq A$.

[1] The proposition (14) has been removed.

(22) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, and F be a subset of the product prebasis for J. Suppose that for every finite subset G of F holds $\Omega_{\prod J} \not\subseteq \bigcup G$. Let x_1 be an element of $J(i)$ and G be a finite subset of F. Suppose $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \subseteq \bigcup G$. Then there exists a subset A_1 of $J(i)$ such that $A_1 \neq \Omega_{J(i)}$ and $x_1 \in A_1$ and $(\mathrm{proj}(J,i))^{-1}(A_1) \in G$ and A_1 is open.

(23) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I, i be an element of I, and F be a subset of the product prebasis for J. Suppose for every element i of I holds $J(i)$ is compact and for every finite subset G of F holds $\Omega_{\prod J} \not\subseteq \bigcup G$. Then there exists an element x_1 of $J(i)$ such that for every finite subset G of F holds $(\mathrm{proj}(J,i))^{-1}(\{x_1\}) \not\subseteq \bigcup G$.

(24) Let I be a non empty set and J be a topological space yielding nonempty many sorted set indexed by I. If for every element i of I holds $J(i)$ is compact, then $\prod J$ is compact.

REFERENCES

[1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.

[2] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.

[3] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/funct_7.html.

[4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[5] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[6] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/compts_1.html.

[7] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_2.html.

[8] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[9] Mariusz Giero. More on products of many sorted algebras. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/pralg_3.html.

[10] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.

[11] Jarosław Gryko. Injective spaces. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/waybel18.html.

[12] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.

[13] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.

[14] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/cantor_1.html.

[15] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[16] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pboole.html.

[17] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[18] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received May 23, 2000

Published January 2, 2004