

N -Dimensional Binary Vector Spaces

Kenichi Arai¹
Tokyo University of Science
Chiba, Japan

Hiroyuki Okazaki
Shinshu University
Nagano, Japan

Summary. The binary set $\{0, 1\}$ together with modulo-2 addition and multiplication is called a binary field, which is denoted by \mathbb{F}_2 . The binary field \mathbb{F}_2 is defined in [1]. A vector space over \mathbb{F}_2 is called a binary vector space. The set of all binary vectors of length n forms an n -dimensional vector space V_n over \mathbb{F}_2 . Binary fields and n -dimensional binary vector spaces play an important role in practical computer science, for example, coding theory [15] and cryptology. In cryptology, binary fields and n -dimensional binary vector spaces are very important in proving the security of cryptographic systems [13]. In this article we define the n -dimensional binary vector space V_n . Moreover, we formalize some facts about the n -dimensional binary vector space V_n .

MSC: 15A03 03B35

Keywords: formalization of binary vector space

MML identifier: NBVECTSP, version: 8.1.01 5.13.1174

The notation and terminology used in this paper have been introduced in the following articles: [6], [1], [2], [16], [5], [7], [11], [17], [8], [9], [18], [24], [14], [4], [25], [26], [19], [23], [12], [20], [21], [22], [27], and [10].

In this paper m, n, s denote non zero elements of \mathbb{N} .

Now we state the proposition:

- (1) Let us consider elements u_1, v_1, w_1 of $Boolean^n$. Then $\text{Op-XOR}((\text{Op-XOR}(u_1, v_1)), w_1) = \text{Op-XOR}(u_1, (\text{Op-XOR}(v_1, w_1)))$.

Let n be a non zero element of \mathbb{N} . The functor $\text{XOR}_B(n)$ yielding a binary operation on $Boolean^n$ is defined by

- (Def. 1) Let us consider elements x, y of $Boolean^n$. Then $it(x, y) = \text{Op-XOR}(x, y)$.

The functor $\text{Zero}_B(n)$ yielding an element of $Boolean^n$ is defined by the term

- (Def. 2) $n \mapsto 0$.

¹This research was presented during the 2013 International Conference on Foundations of Computer Science FCS'13 in Las Vegas, USA.

The functor n -binary additive group yielding a strict additive loop structure is defined by the term

(Def. 3) $\langle \text{Boolean}^n, \text{XOR}_B(n), \text{Zero}_B(n) \rangle$.

Let us consider an element u_1 of Boolean^n . Now we state the propositions:

- (2) $\text{Op-XOR}(u_1, \text{Zero}_B(n)) = u_1$.
- (3) $\text{Op-XOR}(u_1, u_1) = \text{Zero}_B(n)$.

Let n be a non zero element of \mathbb{N} . Note that n -binary additive group is add-associative right zeroed right complementable Abelian and non empty and every element of \mathbf{Z}_2 is Boolean.

Let u, v be elements of \mathbf{Z}_2 . We identify $u \oplus v$ with $u + v$. We identify $u \wedge v$ with $u \cdot v$. Let n be a non zero element of \mathbb{N} . The functor $\text{MLT}_B(n)$ yielding a function from (the carrier of \mathbf{Z}_2) $\times \text{Boolean}^n$ into Boolean^n is defined by

(Def. 4) Let us consider an element a of Boolean , an element x of Boolean^n , and a set i . If $i \in \text{Seg } n$, then $it(a, x)(i) = a \wedge x(i)$.

The functor n -binary vector space yielding a vector space over \mathbf{Z}_2 is defined by the term

(Def. 5) $\langle \text{Boolean}^n, \text{XOR}_B(n), \text{Zero}_B(n), \text{MLT}_B(n) \rangle$.

Let us note that n -binary vector space is finite.

Let us note that every subspace of n -binary vector space is finite.

Now we state the propositions:

- (4) Let us consider a natural number n . Then $\sum n \mapsto 0_{\mathbf{Z}_2} = 0_{\mathbf{Z}_2}$.
- (5) Let us consider a finite sequence x of elements of \mathbf{Z}_2 , an element v of \mathbf{Z}_2 , and a natural number j . Suppose
 - (i) $\text{len } x = m$, and
 - (ii) $j \in \text{Seg } m$, and
 - (iii) for every natural number i such that $i \in \text{Seg } m$ holds if $i = j$, then $x(i) = v$ and if $i \neq j$, then $x(i) = 0_{\mathbf{Z}_2}$.

Then $\sum x = v$. The theorem is a consequence of (4). PROOF: Define $\mathcal{P}[\text{natural number}] \equiv$ for every non zero element m of \mathbb{N} for every finite sequence x of elements of \mathbf{Z}_2 for every element v of \mathbf{Z}_2 for every natural number j such that $\$1 = m$ and $\text{len } x = m$ and $j \in \text{Seg } m$ and for every natural number i such that $i \in \text{Seg } m$ holds if $i = j$, then $x(i) = v$ and if $i \neq j$, then $x(i) = 0_{\mathbf{Z}_2}$ holds $\sum x = v$. For every natural number k such that $\mathcal{P}[k]$ holds $\mathcal{P}[k+1]$ by [3, (11)], [5, (59), (5), (1)]. For every natural number k , $\mathcal{P}[k]$ from [3, Sch. 2]. \square

- (6) Let us consider a (the carrier of n -binary vector space)-valued finite sequence L and a natural number j . Suppose
 - (i) $\text{len } L = m$, and
 - (ii) $m \leq n$, and

(iii) $j \in \text{Seg } n$.

Then there exists a finite sequence x of elements of \mathbf{Z}_2 such that

(iv) $\text{len } x = m$, and

(v) for every natural number i such that $i \in \text{Seg } m$ there exists an element K of Boolean^n such that $K = L(i)$ and $x(i) = K(j)$.

PROOF: Define $\mathcal{Q}[\text{natural number, set}] \equiv$ there exists an element K of Boolean^n such that $K = L(\$_1)$ and $\$_2 = K(j)$. For every natural number i such that $i \in \text{Seg } m$ there exists an element y of Boolean such that $\mathcal{Q}[i, y]$. Consider x being a finite sequence of elements of Boolean such that $\text{dom } x = \text{Seg } m$ and for every natural number i such that $i \in \text{Seg } m$ holds $\mathcal{Q}[i, x(i)]$ from [5, Sch. 5]. \square

(7) Let us consider a (the carrier of n -binary vector space)-valued finite sequence L , an element S of Boolean^n , and a natural number j . Suppose

(i) $\text{len } L = m$, and

(ii) $m \leq n$, and

(iii) $S = \sum L$, and

(iv) $j \in \text{Seg } n$.

Then there exists a finite sequence x of elements of \mathbf{Z}_2 such that

(v) $\text{len } x = m$, and

(vi) $S(j) = \sum x$, and

(vii) for every natural number i such that $i \in \text{Seg } m$ there exists an element K of Boolean^n such that $K = L(i)$ and $x(i) = K(j)$.

The theorem is a consequence of (6). PROOF: Consider x being a finite sequence of elements of \mathbf{Z}_2 such that $\text{len } x = m$ and for every natural number i such that $i \in \text{Seg } m$ there exists an element K of Boolean^n such that $K = L(i)$ and $x(i) = K(j)$. Consider f being a function from \mathbb{N} into n -binary vector space such that $\sum L = f(\text{len } L)$ and $f(0) = 0_{n\text{-binary vector space}}$ and for every natural number j and for every element v of n -binary vector space such that $j < \text{len } L$ and $v = L(j + 1)$ holds $f(j + 1) = f(j) + v$. Define $\mathcal{Q}[\text{natural number, set}] \equiv$ there exists an element K of Boolean^n such that $K = f(\$_1)$ and $\$_2 = K(j)$. For every element i of \mathbb{N} , there exists an element y of the carrier of \mathbf{Z}_2 such that $\mathcal{Q}[i, y]$ by [1, (3)]. Consider g being a function from \mathbb{N} into \mathbf{Z}_2 such that for every element i of \mathbb{N} , $\mathcal{Q}[i, g(i)]$ from [9, Sch. 3]. Set $S_j = S(j)$. $S_j = g(\text{len } x)$. $g(0) = 0_{\mathbf{Z}_2}$ by [1, (5)]. For every natural number k and for every element v_2 of \mathbf{Z}_2 such that $k < \text{len } x$ and $v_2 = x(k + 1)$ holds $g(k + 1) = g(k) + v_2$ by [3, (11), (13)]. \square

(8) Suppose $m \leq n$. Then there exists a finite sequence A of elements of Boolean^n such that

- (i) $\text{len } A = m$, and
- (ii) A is one-to-one, and
- (iii) $\overline{\text{rng } A} = m$, and
- (iv) for every natural numbers i, j such that $i \in \text{Seg } m$ and $j \in \text{Seg } n$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$.

PROOF: Define $\mathcal{P}[\text{natural number, function}] \equiv$ for every natural number j such that $j \in \text{Seg } n$ holds if $\$1 = j$, then $\$2(j) = \text{true}$ and if $\$1 \neq j$, then $\$2(j) = \text{false}$. For every natural number k such that $k \in \text{Seg } m$ there exists an element x of Boolean^n such that $\mathcal{P}[k, x]$. Consider A being a finite sequence of elements of Boolean^n such that $\text{dom } A = \text{Seg } m$ and for every natural number k such that $k \in \text{Seg } m$ holds $\mathcal{P}[k, A(k)]$ from [5, Sch. 5]. For every elements x, y such that $x, y \in \text{dom } A$ and $A(x) = A(y)$ holds $x = y$ by [5, (5)]. \square

- (9) Let us consider a finite sequence A of elements of Boolean^n , a finite subset B of n -binary vector space, a linear combination l of B , and an element S of Boolean^n . Suppose

- (i) $\text{rng } A = B$, and
- (ii) $m \leq n$, and
- (iii) $\text{len } A = m$, and
- (iv) $S = \sum l$, and
- (v) A is one-to-one, and
- (vi) for every natural numbers i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } m$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$.

Let us consider a natural number j . If $j \in \text{Seg } m$, then $S(j) = l(A(j))$. The theorem is a consequence of (7) and (5). PROOF: Set $V = n$ -binary vector space. Reconsider $F_1 = A$ as a finite sequence of elements of V . Consider x being a finite sequence of elements of \mathbf{Z}_2 such that $\text{len } x = m$ and $S(j) = \sum x$ and for every natural number i such that $i \in \text{Seg } m$ there exists an element K of Boolean^n such that $K = (l \cdot F_1)(i)$ and $x(i) = K(j)$. For every natural number i such that $i \in \text{Seg } m$ holds if $i = j$, then $x(i) = l(A(j))$ and if $i \neq j$, then $x(i) = 0_{\mathbf{Z}_2}$ by [5, (5)], [1, (3), (5)]. \square

- (10) Let us consider a finite sequence A of elements of Boolean^n and a finite subset B of n -binary vector space. Suppose

- (i) $\text{rng } A = B$, and
- (ii) $m \leq n$, and
- (iii) $\text{len } A = m$, and
- (iv) A is one-to-one, and

- (v) for every natural numbers i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } m$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$.

Then B is linearly independent. The theorem is a consequence of (9).

PROOF: Set $V = n$ -binary vector space. For every linear combination l of B such that $\sum l = 0_V$ holds the support of $l = \emptyset$ by [1, (5)]. \square

- (11) Let us consider a finite sequence A of elements of Boolean^n , a finite subset B of n -binary vector space, and an element v of Boolean^n . Suppose

- (i) $\text{rng } A = B$, and
- (ii) $\text{len } A = n$, and
- (iii) A is one-to-one.

Then there exists a linear combination l of B such that for every natural number j such that $j \in \text{Seg } n$ holds $v(j) = l(A(j))$. PROOF: Set $V = n$ -binary vector space. Define $\mathcal{Q}[\text{element}, \text{element}] \equiv$ there exists a natural number j such that $j \in \text{Seg } n$ and $\$1 = A(j)$ and $\$2 = v(j)$. For every element x such that $x \in B$ there exists an element y such that $y \in$ the carrier of \mathbf{Z}_2 and $\mathcal{Q}[x, y]$ by [1, (3)]. Consider l_1 being a function from B into the carrier of \mathbf{Z}_2 such that for every element x such that $x \in B$ holds $\mathcal{Q}[x, l_1(x)]$ from [9, Sch. 1]. For every natural number j such that $j \in \text{Seg } n$ holds $l_1(A(j)) = v(j)$ by [8, (3)]. Set $f =$ (the carrier of V) $\mapsto 0_{\mathbf{Z}_2}$. Set $l = f + \cdot l_1$. For every element v of V such that $v \notin B$ holds $l(v) = 0_{\mathbf{Z}_2}$ by [17, (7)]. For every element x such that $x \in$ the support of l holds $x \in B$. For every natural number j such that $j \in \text{Seg } n$ holds $v(j) = l(A(j))$ by [8, (3)]. \square

- (12) Let us consider a finite sequence A of elements of Boolean^n and a finite subset B of n -binary vector space. Suppose

- (i) $\text{rng } A = B$, and
- (ii) $\text{len } A = n$, and
- (iii) A is one-to-one, and
- (iv) for every natural numbers i, j such that $i, j \in \text{Seg } n$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$.

Then $\text{Lin}(B) =$ (the carrier of n -binary vector space, the addition of n -binary vector space, the zero of n -binary vector space, the left multiplication of n -binary vector space). The theorem is a consequence of (11) and (9).

PROOF: Set $V = n$ -binary vector space. For every element $x, x \in$ the carrier of $\text{Lin}(B)$ iff $x \in$ the carrier of V by [5, (13)], [22, (7)]. \square

- (13) There exists a finite subset B of n -binary vector space such that

- (i) B is a basis of n -binary vector space, and
- (ii) $\overline{\overline{B}} = n$, and

- (iii) there exists a finite sequence A of elements of $Boolean^n$ such that $\text{len } A = n$ and A is one-to-one and $\overline{\text{rng } A} = n$ and $\text{rng } A = B$ and for every natural numbers i, j such that $i, j \in \text{Seg } n$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$.

The theorem is a consequence of (8), (10), and (12).

- (14) (i) n -binary vector space is finite dimensional, and
 (ii) $\dim(n\text{-binary vector space}) = n$.

The theorem is a consequence of (13).

Let n be a non zero element of \mathbb{N} . One can verify that n -binary vector space is finite dimensional.

Now we state the proposition:

- (15) Let us consider a finite sequence A of elements of $Boolean^n$ and a subset C of n -binary vector space. Suppose
- (i) $\text{len } A = n$, and
 - (ii) A is one-to-one, and
 - (iii) $\overline{\text{rng } A} = n$, and
 - (iv) for every natural numbers i, j such that $i, j \in \text{Seg } n$ holds if $i = j$, then $A(i)(j) = \text{true}$ and if $i \neq j$, then $A(i)(j) = \text{false}$, and
 - (v) $C \subseteq \text{rng } A$.

Then

- (vi) $\text{Lin}(C)$ is a subspace of n -binary vector space, and
- (vii) C is a basis of $\text{Lin}(C)$, and
- (viii) $\dim(\text{Lin}(C)) = \overline{C}$.

The theorem is a consequence of (10).

REFERENCES

- [1] Jesse Alama. The vector space of subsets of a set based on symmetric difference. *Formalized Mathematics*, 16(1):1–5, 2008. doi:10.2478/v10037-008-0001-7.
- [2] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [3] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [4] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [6] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [7] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [8] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [9] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.

- [10] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [11] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [12] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [13] X. Lai. Higher order derivatives and differential cryptanalysis. *Communications and Cryptography*, pages 227–233, 1994.
- [14] Robert Milewski. Associated matrix of linear map. *Formalized Mathematics*, 5(3):339–345, 1996.
- [15] J.C. Moreira and P.G. Farrell. *Essentials of Error-Control Coding*. John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, 2006.
- [16] Hiroyuki Okazaki and Yasunari Shidama. Formalization of the data encryption standard. *Formalized Mathematics*, 20(2):125–146, 2012. doi:10.2478/v10037-012-0016-y.
- [17] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [18] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [19] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [20] Wojciech A. Trybulec. Subspaces and cosets of subspaces in vector space. *Formalized Mathematics*, 1(5):865–870, 1990.
- [21] Wojciech A. Trybulec. Linear combinations in vector space. *Formalized Mathematics*, 1(5):877–882, 1990.
- [22] Wojciech A. Trybulec. Basis of vector space. *Formalized Mathematics*, 1(5):883–885, 1990.
- [23] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [24] Edmund Woronowicz. Many argument relations. *Formalized Mathematics*, 1(4):733–737, 1990.
- [25] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [26] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [27] Mariusz Żynel. The Steinitz theorem and the dimension of a vector space. *Formalized Mathematics*, 5(3):423–428, 1996.

Received April 17, 2013
